



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

RECEIVED

17 NOV 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03023815.8

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Anmeldung Nr:
Application no.: 03023815.8
Demande no:

Anmeldetag:
Date of filing: 17.10.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

DEUTSCHES KREBSFORSCHUNGSZENTRUM STIFTUNG
DES ÖFFENTLICHEN RECHTS
Im Neuenheimer Feld 280
69120 Heidelberg
ALLEMAGNE
Ruprecht-Karls-Universität Heidelberg
Grabengasse 1
69117 Heidelberg
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Use of ADAM 12 for diagnosis and therapy of preeclampsia

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G01N33/48

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

21. Okt. 2003

- 1 -

Deutsches Krebsforschungszentrum
Universität HeidelbergOctober 17, 2003
DK62208EP HN/FLZ/ROE/hen/hmü

5 **Use of ADAM 12 for diagnosis and therapy of preeclampsia**

10 The present invention relates to the use of ADAM 12, particularly ADAM 12-S, for diagnosis and therapy of preeclampsia and related syndromes.

15 Preeclampsia (also known as toxemia or eclampsism) is a disorder or syndrome affecting about 5 % of pregnant women. It becomes manifest in the last trimester of pregnancy and is characterized by high blood pressure, protein in the urine, and – sometimes – swelling (edema) of the face and hands. Preeclampsia may also include sudden weight gain, blurred vision, severe headaches, dizziness, and stomach pain.

20 In preeclampsia, the trophoblast (embryonic tissue that invades the uterus wall to form the embryonic part of the placenta) is poorly differentiated and shows incomplete invasion of the uterus wall.

25 When left untreated, preeclampsia can lead to severe problems, endangering both mother and fetus. High blood pressure constricts the blood vessels in the uterus that supply the fetus with oxygen and nutrients. The fetal growth is retarded. Preeclampsia can increase the risk of placental abruption (separation of the placenta from the uterus wall) before delivery, risk of kidney failure, cerebral hemorrhage, intravascular coagulation, lung edema, and circulatory collapse.

30 Furthermore, untreated preeclampsia can lead to a related life-threatening condition, eclampsia. Eclampsia can lead to convulsions and coma. The HELLP (hemolysis, elevated liver enzymes, and low platelet count) syndrome is also frequently considered to be related to preeclampsia, as preeclampsia is thought to increase the risk for HELLP. HELLP is characterized by destruction of red blood cells, liver damage, and a reduction in the number of blood platelets. Thus, also HELLP can be life-threatening.

Intrauterine growth retardation (small for gestational age (SGA) infants) is another syndrome related to preeclampsia, as it is associated with poor trophoblast differentiation and growth retardation of the embryo.

Pregnancy induced hypertension (PIH) is also a syndrome related to preeclampsia. It is characterized by high blood pressure, but lacks the proteinuria characteristic for preeclampsia.

Other syndromes related to preeclampsia are superimposed gestosis ("Aufpropfgestose") and gestational diabetes.

A causal treatment of preeclampsia and said related syndromes is not available. The usual treatment consists of the treatment of hypertension, intravenous magnesium to prevent eclamptic convulsions and corticosteroids in case of HELLP syndrome. Very often symptoms can not be controlled and preterm delivery is the only way to save the life of mother and child. Preterm delivery is still associated with a high morbidity and mortality of the child.

Currently, diagnosis of preeclampsia is limited to relatively late arising symptoms such as blood pressure (typically above 140/90 mm Hg) and protein content of the urine (proteinuria, typically more than 0.3 g/24 hours or a level of 2+ by dipstick test). Another sign for preeclampsia is glomerular endotheliosis, a histological change of the kidney which can be observed in kidney biopsies of 50% of the patients. These markers are relatively unreliable or become manifest only at later stages of preeclampsia. Similarly, said related syndromes can not be diagnosed in a satisfactory manner. In summary, there remains an urgent need for molecular markers indicative of preeclampsia (Beaulieu, M.D., Prevention of preeclampsia. In: Canadian Task Force on the Periodic Health Examination. Canadian Guide to Clinical Preventive Health Care. Ottawa: Health Canada, 1994; 136-43).

A small number of genes has been reported to be either upregulated (EPAS-1/HIF-2 α , neurokinin B, TIMP-1, VEGFR-1 (=sFLT-1), VEGF, IGFBP-1, IGFBP-3, matrix metalloproteinase-2, leptin, plasminogen activator inhibitor 1 (PAI-1)) or downregulated (IGF-1, angiopoietin-2, decorin, various ephrins, PIGF, HLA-G) during preeclampsia. PIGF has been reported to be downregulated in blood serum, but not in placental tissue. HB-EGF, various ephrins, and TGF- β 3 have also been discussed as being de-regulated during preeclampsia.

However, most of these genes are only weakly up- or downregulated and/or can only be measured in placental tissue, requiring to take placenta tissue samples for analysis. Thus, frequent measurements at short time intervals are not possible. In any case, none of the genes cited above have been put into practice as routine markers for preeclampsia or related syndromes.

Consequently, a reliable diagnosis of preeclampsia or related syndromes is not possible at the moment. However, reliable diagnosis of onset and progression of said disorders or syndromes would allow to develop better treatment and to evaluate the risk for mother and fetus at each stage of pregnancy. Reliable diagnosis is also needed, because women experience no obvious symptoms at early stages of preeclampsia or related syndromes.

Therefore, there is a persistent need for molecular markers of preeclampsia or related syndromes. Such markers would also be useful to identify and classify different subtypes of preeclampsia and related syndromes.

Consequently, the problem underlying the present invention resides in improving diagnosis of preeclampsia or related syndromes. In particular, the problem resides in providing a marker more indicative of preeclampsia and/or related syndromes than other markers. In particular, the problem resides in providing a highly indicative marker that can also be measured in body fluids.

Another problem underlying the present invention resides in providing treatment of preeclampsia and/or related syndromes.

With respect to diagnosis, the problem is solved by use of a nucleic acid molecule comprising a nucleic acid selected from the group consisting of

- a) a nucleic acid with a sequence as presented in SEQ ID NO: 1 or 3;
- b) a nucleic acid with a sequence that exhibits a sequence identity with any of the sequences defined in a) of at least 70% over 300 residues;
- c) a nucleic acid which is capable of hybridizing with the nucleic acid as defined in a), or b) under conditions of medium or high stringency;
- d) a nucleic acid with the antisense-sequence of any of the sequences defined in a), b) or c);
- e) a fragment of any of the nucleic acids as defined in a), b), c), or d);

f) an RNA corresponding to any of the sequences defined in a), b), c), d) or e);

for diagnosis of preeclampsia or a related syndrome.

5 The problem is also solved by use of a ligand binding specifically to a peptide or polypeptide with a sequence selected from the group consisting of

- a) an amino acid sequence as presented in SEQ ID NO: 2 or 4;
- b) an amino acid sequence exhibiting a sequence identity with any of the
- 10 amino acid sequences according to a) of at least 85% over 100 amino acid residues;
- c) a fragment of any of the sequences defined above

for diagnosis of preeclampsia or a related syndrome.

15

For reference, the individual SEQ ID NO denotes the following sequence:

SEQ ID NO: 1	nucleotide sequence of the human ADAM 12-L cDNA
SEQ ID NO: 2	protein sequence of the human ADAM 12-L protein
20 SEQ ID NO: 3	nucleotide sequence of the human ADAM 12-S cDNA
SEQ ID NO: 4	protein sequence of the human ADAM 12-S protein
SEQ ID NO: 5	nucleotide sequence region specific for the human ADAM 12-L cDNA
SEQ ID NO: 6	protein sequence region specific for human ADAM 12-L
25 SEQ ID NO: 7	nucleotide sequence region specific for the human ADAM 12-S cDNA
SEQ ID NO: 8	protein sequence region specific for human ADAM 12-S
SEQ ID NO: 9	PCR primer (1), ADAM 12 5' primer
SEQ ID NO: 10	PCR primer (2), ADAM 12-L 3' primer
30 SEQ ID NO: 11	PCR primer (3), ADAM 12-L 5' primer
SEQ ID NO: 12	PCR primer (4), ADAM 12-L 3' primer
SEQ ID NO: 13	PCR primer (5), ADAM 12-S 3' primer, containing a single nucleotide exchange (G instead of C) at position 15.
SEQ ID NO: 14	PCR primer (6), ADAM 12-S 3' primer, correct sequence
35 SEQ ID NO: 15	PCR primer (7), ADAM 12-S 5' primer
SEQ ID NO: 16	PCR primer (8), ADAM 12-S 3' primer
SEQ ID NO: 17	PCR primer (9), human cyclophilin B 5' primer

SEQ ID NO: 18 PCR primer (10), human cyclophilin B 3' primer
SEQ ID NO: 19 PCR primer (11), EF-1 α 5' primer
SEQ ID NO: 20 PCR primer (12), EF-1 α 3' primer

5

SEQ ID NO: 1 to 16 refer to nucleic acids, peptides or polypeptides derived from ADAM 12. ADAM 12 (meltrin alpha) is a member of the ADAM (a disintegrin and metalloprotease) family. Human ADAM 12 exists in two isoforms, a longer membrane anchored isoform, ADAM 12-L (SEQ ID NO: 1, 2, GenBank Acc. No. AF023476) and a
10 shorter secreted isoform, ADAM 12-S (SEQ ID NO: 3, 4, GenBank Acc. No. AF023477).

ADAM 12 has been implicated in cell adhesion and muscle cell differentiation and fusion. Additionally, ADAM 12 has been implicated in fat tissue development and it is upregulated in several types of human carcinomas. ADAM 12-S is expressed highly in the
15 placenta and can be detected in blood serum during pregnancy, while it is undetectable in non-pregnancy serum. However, no connection of ADAM 12, particularly ADAM 12-S, to preeclampsia or a related syndrome has been reported previously.

As a consequence of the present invention it is known that expression, particularly
20 transcription, of ADAM 12 is strongly upregulated in preeclampsia. In a comparison with 1600 known genes examined by gene array analysis, ADAM 12 was found to be the most strongly upregulated gene (Example 8). Therefore, ADAM 12 can be considered a valuable diagnostic marker, and upregulation of ADAM 12 can be used to diagnose preeclampsia and related syndromes. ADAM 12, particularly ADAM 12-S, is present in body fluids such
25 as blood. Blood samples are taken routinely at physical examinations during most pregnancies. Thus, it is possible to routinely monitor pregnant mothers at short intervals for upregulation of ADAM 12, allowing to early detect and monitor preeclampsia or related disorders with minimal effort and risk for the mother.

30 "Nucleic acid" according to the present invention relates to all known nucleic acids such as DNA, RNA, peptide nucleic acids, morpholinos, and nucleic acids with backbones other than phosphodiester, such as phosphothioates or phosphoramidates. The person skilled in the art knows which nucleic acids are best suited for a particular purpose, such as diagnosis, or a particular method (such as Northern hybridization or polymerase chain
35 reaction (PCR)).

Unless otherwise specified, the manipulations of nucleic acids and polypeptides/-proteins can be performed using standard methods of molecular biology and immunology (see, e.g. Maniatis et al. (1989), *Molecular cloning: A laboratory manual*, Cold Spring Harbor Lab., Cold Spring Harbor, NY; Ausubel, F.M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Tijssen, P., *Practice and Theory of Enzyme Immunoassays*, Elsevier Press, Amsterdam, Oxford, New York, 1985).

The term "comprise" preferably refers to nucleic acid molecules in which the nucleic acids with the described sequences are functionally relevant, e.g. for diagnostic use or therapeutic use, such as vectors for therapeutical use or expression of corresponding RNAs or proteins. For example, the above defined nucleic acids may be part of a vector allowing transcription of antisense DNA or RNA probes. Preferably, any additional nucleic acids upstream or downstream of the nucleic acid are not longer than 80 kb, more preferably not longer than 20kb, more preferably not longer than 5 kb. More preferably, the term "comprise" does not relate to large constructs accidentally including the sequence, such as genomic BAC or YAC clones. More preferably, the nucleic acid molecule consists of the above defined nucleic acids.

Preferably, the nucleic acid molecule is purified and/or isolated. Suitable purification methods, including gel electrophoresis and capillary electrophoresis, are well known in the art. An isolated nucleic acid molecule is understood as a nucleic acid molecule isolated from the human or animal body or gained by means of a technical process, such as a microbiological process or organic synthesis.

In a preferred embodiment, said nucleic acid molecule comprises a nucleic acid with a sequence selected from the group of sequences as presented in SEQ ID NO: 1 or 3. SEQ ID NO: 1 corresponds to the sequence presented in GenBank Acc. No. AF023476, and SEQ ID NO: 3 corresponds to the sequence presented in GenBank Acc. No. AF023477.

In another preferred embodiment, said nucleic acid molecule comprises a nucleic acid with a sequence that exhibits a sequence identity with any of the sequences defined above of at least 70%, preferably at least 80%, more preferably at least 90%, and most preferred of at least 95% over 300 nucleotide residues.

"Sequence identity" refers to the degree of identity (% identity) of two sequences that in the in the case of peptides or polypeptides can be determined by means of for example BLASTP 2.0.1 and in the case of nucleic acids can be determined by means of for example BLASTN 1.014, wherein the Filter is set off and BLOSUM is 62 (Altschul et al., Nucleic
5 Acids Res., 25:3389-3402).

Such nucleic acids include those whose sequences deviate from the above defined sequences due to the degeneration of the genetic code, the presence of different alleles in the human or animal population, and the presence of related sequences in different species.
10 Such nucleic acids can still be used according to the present invention.

In another preferred embodiment, the nucleic acid molecule comprises a nucleic acid capable of hybridizing with the nucleic acids according to SEQ ID NO: 1 or 3 under conditions of low, preferably medium, or, more preferred, high stringency.

15 In such hybrids, duplex formation and stability depend on substantial complementarity between the two strands of the hybrid and a certain degree of mismatch can be tolerated. Therefore, the nucleic acids and nucleic acid probes of the present invention may include mutations (both single and multiple), deletions, insertions of the above-identified
20 sequences, and combination thereof, as long as that nucleic acids still have substantial sequence similarity to the original sequence which permits the formation of stable hybrids with the target nucleotide sequence of interest. For example, a single nucleotide mismatch in PCR primer (SEQ ID NO: 13) as compared to the correct primer sequence (SEQ ID NO: 14) was found to be tolerable.

25 Suitable experimental conditions for determining whether a given DNA or RNA sequence "hybridizes" to a specified polynucleotide or oligonucleotide probe involve presoaking of the filter containing the DNA or RNA to examine for hybridization in 5 x SSC (sodium chloride/sodium citrate) buffer for 10 minutes, and prehybridization of the filter in a
30 solution of 5 x SSC, 5 x Denhardt's solution, 0,5 % SDS and 100 mg/ml of denaturated sonicated salmon sperm DNA (Maniatis et al., 1989, see above), followed by hybridization in the same solution containing a concentration of 10 ng/ml of a random primed (Feinberg, A.P. and Vogelstein, B. (1983), Anal. Biochem. 132:6-13), ³²P-dCTP-labeled (specific activity > 1 x 10⁹ cpm/μg) probe for 12 hours at approximately 45°C. The filter is then
35 washed twice for 30 minutes in 2 x SSC, 0,5% SDS at at least 55°C (low stringency), at

least 60°C (medium stringency), preferably at least 65°C (medium/high stringency), more preferably at least 70°C (high stringency) or most preferably at least 75°C (very high stringency). Molecules to which the probe hybridizes under the chosen conditions are detected using an x-ray film or a "phosphor imager".

5

In another preferred embodiment, said nucleic acid molecule comprises a nucleic acid which is capable of hybridizing with the nucleic acids according to SEQ ID NO: 1 or 3, but not capable of hybridizing with nucleic acids encoding ADAM 19 or ADAM 33.

- 10 Preferably, the nucleic acids should be chosen from sequence regions specific for the above-defined sequences, i.e. the nucleic acids should not cross-hybridize with nucleic acids encoding other genes, particularly other metalloprotease genes, more particularly ADAMs (Seals, D.F. and Courtneidge, S.A. (2003). The ADAMs family of metalloproteases: multidomain proteins with multiple functions. *Genes & Dev.* 17, 7-30),
15 even more particularly ADAM 19 (GenBank Acc. No. Y13786) or ADAM 33 (GenBank Acc. No. AB009672, most particularly ADAM 2 (fertilin beta, GenBank Acc. Nos. AJ133005 or NM_001464) (see also Yoshinaka, T., Nishii, K., Yamada, K., Sawada, H., Nishiwaki, E., Smith, K., Yoshino, K., Ishiguro, H., and Higashiyama, S. (2002). Identification and characterization of novel mouse and human ADAM 33s with potential
20 metalloprotease activity. *Gene* 282, 227-236), under the applied conditions, preferably conditions of medium, medium/high, high or very high stringency.

In another preferred embodiment, the nucleic acid molecule comprises a nucleic acid with the antisense-sequence of any of the sequences defined above.

25

In another preferred embodiment, the nucleic acid molecule comprises a nucleic acid which is a fragment of any of the nucleic acids as defined above.

- 30 The term "fragment" as used according to the present invention can have different meanings depending on the molecule and purpose referred to. A person skilled in the art knows how to choose appropriate fragments for the relevant purpose. In particular, the person skilled in the art knows how to choose fragments of appropriate length and suitable sequence (see also Examples 9 and 10).

- 35 Preferably, the fragments should be chosen from sequence regions specific for the above-defined sequences, i.e. the chosen fragments should not cross-hybridize with nucleic acids

encoding other genes, particularly other metalloprotease genes, more particularly ADAMs, even more particularly ADAM 19 or ADAM 33, most particularly ADAM 2, under the applied conditions. Sequence regions specific for the above defined sequences can be determined easily by the person skilled in the art, e.g. by database searches and sequence
5 comparisons. Such searches and comparisons are routine in the art and possible using sequence analysis programs such as BLAST and ClustalW. Preferably, the fragments should be chosen from regions of high complexity which can be determined by low complexity filters available in standard sequence analysis programs. Specificity for the purpose of the present invention may also be achieved in spite of cross-hybridization, if the
10 investigated sequences can still be measured unequivocally, e.g. by the size of amplified PCR products. Therefore, such fragments, particularly PCR primers, can still be used according to the present invention.

In another preferred embodiment, the fragment has a sequence consisting of or being part
15 of any of SEQ ID NO: 5, 7, 9, 10, 11, 12, 13, 14, 15, 16;

In another preferred embodiment, the fragment should be at least 12, preferably at least 15, more preferably at least 19, even more preferably at least 30 nucleotides in length.

20 In another preferred embodiment, said nucleic acid molecule comprises an RNA corresponding to any of the sequences defined above. RNA molecules are frequently used as probes for analysis of nucleic acids and measurement of gene expression, e.g. by in situ hybridization or RNase protection assays. A suitable RNA in situ hybridization probe may, for example, comprise or consist of the 667 nucleotide sequence fragment
25 corresponding to the antisense sequence of position 1769 to 2445 in SEQ ID NO: 3.

Diagnosis according to the present invention includes monitoring, confirmation, subclassification and prediction of the relevant disorder or syndrome. Monitoring relates to monitoring of an already diagnosed disorder, e.g. to analyze the progression of the disorder
30 or the influence of a particular treatment on the progression of disorder. Confirmation relates to the confirmation or substantiation of a diagnosis already performed using other indicators, symptoms, or markers. Subclassification relates to further defining a diagnosis according to different subclasses of the diagnosed disorder, e.g. defining according to mild and severe forms of the disorder. Prediction relates to predicting a disorder before other
35 symptoms or markers have become evident or have become significantly altered. Prediction also relates to classifying patients into groups of low, medium, or high risk of developing the disorder.

In another preferred embodiment, diagnosis relates to the diagnosis of syndromes related to preeclampsia.

5 According to the present invention, syndromes "related" to preeclampsia include other pregnancy-associated disorders or syndromes with similar aetiology and/or symptoms as preeclampsia. Particularly included are eclampsia, pregnancy induced hypertension, HELLP syndrome, intrauterine growth retardation and superimposed gestosis. Most particularly, related syndromes are eclampsia, pregnancy induced hypertension, HELLP
10 syndrome, and superimposed gestosis.

For diagnosis of preeclampsia or related syndromes, preferably the expression level or amount of ADAM 12 RNA or its corresponding protein is determined. The term "measuring" according to the present invention relates to determining the amount or
15 activity, preferably semi-quantitatively or quantitatively, of the substance in question. Measuring is frequently done indirectly, through measuring of ligands, labels, or enzymatic reaction products.

ADAM 12 expression can be measured in tissue samples and body fluid samples. A tissue
20 sample refers to any kind of tissue obtained from the dead or alive human or animal body, particularly a placental tissue sample, more particularly a human placental tissue sample. Body fluid according to the present invention refers to any kind of body fluid, particularly to blood, amniotic fluid, lymph, cerebral liquor, urine, and saliva, more particularly to blood, amniotic fluid, and urine, most particularly to blood. Expression according to the
25 present invention includes transcription, translation and transcript or protein stability.

RNA can be measured easily and quantitatively by methods known to the person skilled in the art. RNA can be obtained or isolated by purification methods known in the art, e.g. involving steps of enzymatic digestion of tissue or proteins, and
30 phenol/chloroform/isoamylalcohol extraction.

Suitable methods for measuring ADAM 12 expression, particularly ADAM 12 RNA (transcript), include the hybridization of nucleic acid probes, such as in situ hybridizations, Northern hybridizations, RNase protection assays, gene arrays, RT-PCR, and other
35 suitable methods known in the art. Preferably, a method allowing quantitative or semi-quantitative measurement, such as quantitative RT-PCR, is chosen.

Characteristics of suitable nucleic acid probes, e.g. in situ hybridization probes or PCR primers, are known in the art. The probes may be based on DNA, RNA or derivatives thereof such as peptide nucleic acids. The probes may be synthesized by use of DNA synthesizers according to standard procedures or, preferably for long sequences, by use of PCR technology with a selected template sequence and selected primers. Depending on the procedure, the nucleic acid probes may be labeled with any suitable substances known in the art, such as digoxigenin, biotin, or thyroxine (as targets of a secondary ligands, particularly a secondary antibody or a fragment thereof), luminescent or fluorescent labels (e.g. fluorescein) and radioactive isotopes (^{32}P , ^{125}I , ^{35}S , and the like). A probe labeled with a radioactive isotope can be constructed from a DNA template by a conventional nick translation reaction or by randomly primed polymerization. The probe may also be labeled at both ends with different types of labels, for example with an isotopic label at one end and a biotin label at the other end. The labeled probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs.

Although gene transcripts are usually present only within cells, ADAM 12 RNA can also be measured (at lower levels) in body fluids, particularly in blood samples, including blood plasma and serum, by methods known in the art (Ng, E.K.O., Tsui, N.B.Y., et al. (2003). Messenger RNA of placental origin is readily detectable in maternal plasma. Proc Natl Acad Sci USA, vol. 100(8):4748-53 and Ng, E.K.O., Leung, T.N., et al. (2003). The concentration of circulating Corticotropin-releasing hormone RNA in maternal plasma is increased in preeclampsia. Clin. Chem. 49, 727-731). According to the present invention, blood plasma is the fluid part of the blood, it lacks the corpuscular components such as blood cells and platelets. Blood serum is blood plasma additionally lacking fibrinogen.

Diagnosis is preferably performed in vitro, i.e. using elements isolated from the body. Tissue samples and body fluid samples can be obtained by laboratory assistants trained for this purpose during routine examinations. Body fluid and tissue samples can be obtained by any method known to the person skilled in the art, including amniocentesis and biopsy (particularly chorion villi biopsy). In particular, blood samples are taken routinely at physical examinations during most pregnancies.

Routine examinations are typically performed in 4 week intervals during the first two trimesters, and biweekly during the last trimester. Typically, blood samples are taken in week 13 and at least once between weeks 21 and 28 of pregnancy. However, many pregnant women come to examinations on a bi-weekly or weekly basis, particularly during the last half of pregnancy.

Thus, it is possible to routinely monitor pregnant women at short intervals for upregulation of ADAM 12, allowing to early detect and monitor preeclampsia or related disorders with minimal effort and risk for mother and fetus.

5

Levels (amounts) of ADAM 12 expression indicative or of diagnostic value for preeclampsia and its related syndromes can be detected by comparing expression levels under healthy conditions with expression levels under sick conditions (conditions of the existing or developing relevant disorder or syndrome). For example, the levels under sick conditions can be determined relative to a healthy control group, or relative the mean population of pregnant mothers, or relative to earlier measurements for the same person.

10

In the present invention, approximately 4 to 5-fold increases of ADAM 12 transcript levels as measured by quantitative Real-Time RT-PCR were found to be correlated with preeclampsia (see Example 10). The transcript levels were increased in placental tissue of both maternal and fetal origin.

15

Further levels of ADAM 12 expression indicative of preeclampsia and related syndromes can be determined in more detail in clinical screenings as known in the art. For example, a level deviating from the average level (e.g. the mean level of the entire investigated group) by more than two standard deviations, preferably by more than three standard deviations may be considered indicative of the analyzed disorder. Deviations indicative of preeclampsia or a related syndrome may be even smaller, if additional markers indicative of the disorder are also deviating from the average level. More weakly deviating levels may be used to classify a person into a medium or high risk group of developing the disorder. Those markers may include blood pressure, protein content in the urine and molecular markers. Additional factors may influence the diagnosis, e.g. the regular blood pressure or the number of previous pregnancies in a given patient. Usually, the highest risk for preeclampsia is during the first pregnancy. Thus, measuring ADAM 12 expression yields a diagnostic parameter that is preferably compared with additional parameters to allow a decision about a necessary treatment.

20

25

30

The present invention also relates to a method of diagnosis of preeclampsia in which the above defined nucleic acid molecules are used.

35

In another preferred embodiment, the fragment has a sequence consisting of or being part of SEQ ID NO: 7, 13, 14, 15, 16.

In another preferred embodiment, the fragment has a sequence consisting of or being part of SEQ ID NO: 5 or 10, 11, 12.

5 SEQ ID NO: 7, 13, 14, 15, and 16 are unique (specific) to the splice variant ADAM 12-S. SEQ ID NO: 5 or 10, 11, 12 are unique (specific) to the splice variant ADAM 12-L. Both splice variants, ADAM 12-L and ADAM 12-S, are upregulated in preeclampsia. However, the splice variants show different regulation properties. In a control group, the level of ADAM 12-L RNA was found to be close to zero, whereas ADAM 12-S showed a
10 measurable basal expression level (both measured by semi-quantitative RT-PCR). Also, the total (upregulated) level of ADAM 12-L RNA in preeclampsia was lower than the one of ADAM 12-S RNA.

For specific detection of mRNA encoding the long splice variant, ADAM 12-L, or the
15 short splice variant, ADAM 12-S, the probe should be directed against the sequence regions unique to these splice variants. For example, probes may be directed against the unique 3'-ends of ADAM 12-L (SEQ ID NO: 5) and ADAM 12-S (SEQ ID NO: 7). As known to the person skilled in the art, a short overlap of the probes with regions common to both splice variants may be tolerable without loss of specificity. For example, the 5'
20 ends of PCR primers may be located in sequence regions common to both transcripts. It is also sufficient if only one of the two primers is located in the regions unique to each splice variant. E.g., the primer according to SEQ ID NO: 9 may be used in combination with the primers according to SEQ ID NO: 13, 14, or 16 to amplify a product specific for ADAM 12-S.

25 As mentioned above, the invention also relates to a the use of a ligand binding a peptide or polypeptide as defined above.

Thus, the present invention also relates to the measurement of ADAM 12 protein. In
30 another preferred embodiment, the biological activity of ADAM 12 is measured. ADAM 12 protein or biological activity can be measured in tissue samples, e.g. in placental tissue samples or other samples taken from the patient.

ADAM 12, particularly ADAM 12-S, is also present in body fluids, particularly the blood
35 (Shi, Z., Xu, W., Loechel, F., Wewers, U.M., and Murphy, L.J. (2000). ADAM 12, a disintegrin metalloprotease, interacts with insulin-like growth factor-binding protein-3. J. Biol. Chem. 275, 18574-18580; Gilpin, B.J., Loechel, F., Mattei, M.-G., Engvall, E.,

Albrechtsen, R., and Wewer, U.M. (1998). A Novel, Secreted Form of human ADAM 12 (Meltrin alpha) Provokes Myogenesis in Vivo. J. Biol. Chem. 273, 157-166). Therefore, ADAM 12-S amount or level of biological activity can easily be measured in body fluids, particularly in blood, blood plasma or blood serum.

5

Biological activity according to the present invention relates to any activity exerted by ADAM 12, ADAM 12-L or ADAM 12-S in the human or animal body. In particular, biological activity relates to proteolytic activity, in particular to proteolytic activity exerted against HB-EGF, IGFBP-3, or IGFBP-5. According to the present invention, the term
10 "biological activity" also relates to any effects, particularly undesired or toxic effects, exerted by ADAM 12, ADAM 12-L, or ADAM 12-S if they are more highly expressed or more active than usual.

In a preferred embodiment, the peptide or polypeptide may comprise a sequence selected
15 from the amino acid sequences presented in SEQ ID NO: 2 or 4. SEQ ID NO: 2 corresponds to the sequence presented in GenBank Acc. No. AF023476 and SEQ ID NO: 4 corresponds to the sequence presented in GenBank Acc. No. AF023477.

In another preferred embodiment, said peptide or polypeptide has a sequence exhibiting a
20 sequence identity with any of the amino acid sequences defined above of at least 85%, preferably 90%, more preferably 95% over 100 amino acid residues.

In another preferred embodiment, said peptide or polypeptide has a sequence which is a
25 fragment of any of the amino acid sequences defined above.

25

The above defined peptide or polypeptide may also be part of fusion peptides or polypeptides. In such fusions another peptide or polypeptide may be fused at the N-terminus or the C-terminus of the peptide or polypeptide of interest. A fusion can be produced by fusing a nucleic acid sequence (or a portion thereof) encoding another
30 polypeptide to a nucleic acid sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art and include ligating the coding sequences so that they are in frame and the expression of the fusion polypeptide is under control of the same promotor(s) and terminator.

35 With respect to diagnosis, the definitions given above for the use of nucleic acids apply analogously.

Suitable methods for measurement of ADAM 12 are known to the person skilled in the art. For example, measurement can be done indirectly, through measuring of cellular responses, the amount of bound ligands, labels, or enzymatic reaction products.

5

For measuring cellular responses, the sample or processed sample is added to a cell culture and an internal or external cellular response is measured. The cellular response may include the expression of a reporter gene or the secretion of a substance, e.g. a peptide, polypeptide, or a small molecule.

10

Other preferred methods for measurement may include measuring the amount of a ligand binding specifically to the ADAM 12 proteins. Binding according to the present invention includes both covalent and non-covalent binding.

15

A ligand according to the present invention can be any peptide, polypeptide, nucleic acid, or other substance (including small molecules) binding to peptides, polypeptides with the above described sequences. It is well known that the above-defined peptides or polypeptides, if obtained or purified from the human or animal body, can be modified, e.g. by glycosylation. A suitable ligand according to the present invention may bind these peptides or polypeptides also via such sites.

20

Preferably, the ligand should bind specifically to said peptides or polypeptides. "Specific binding" according to the present invention means that the ligand should not bind substantially to ("cross-react" with) another protein, isoform or substance present in the sample investigated. Preferably, the specifically bound protein or isoform should be bound with at least 3 times higher, more preferably at least 10 times higher and even more preferably at least 50 times higher affinity than any other relevant protein, isoform, or substance. Other relevant proteins may include other ADAMs, particularly ADAM 2, more particularly ADAM 19 or ADAM 33.

25

30

Non-specific binding may be tolerable, particularly if the investigated ADAM 12 proteins or isoforms can still be distinguished and measured unequivocally, e.g. according to their size on a Western Blot, or by their relatively higher abundance in the sample.

Binding of the ligand can be measured by any method known in the art. Preferably, the method is semi-quantitative or quantitative. Suitable methods are described in the following.

- 5 If the ligand does not only bind, but also serves as a substrate of the above-defined peptides or polypeptides, the reaction product may be measured directly (e.g. cleaving of HB-EGF, IGFBP-3, or IGFBP-5 as a substrate may be measured by analyzing the cleaved product, e.g. on a Western Blot or by ELISA).
- 10 Furthermore, the ligand may be coupled covalently or non-covalently to a label allowing detection and measurement of the ligand.

Labeling may be done by direct or indirect methods. Direct labeling involves coupling of the label directly (covalently or non-covalently) to the ligand. Indirect labeling involves
15 binding (covalently or non-covalently) of a secondary ligand to the first ligand. The secondary ligand should specifically bind to the first ligand. Said secondary ligand may be coupled with a suitable label and/or be the target (receptor) of tertiary ligand binding to the secondary ligand. The use of secondary, tertiary or even higher order ligands is often advantageous to increase the signal. Suitable secondary and higher order ligands may
20 include antibodies, secondary antibodies, and the well-known streptavidin-biotin system (Vector Laboratories, Inc.)

The ligand or substrate may also be "tagged" with one or more tags as known in the art. Such tags may then be targets for higher order ligands. Suitable tags include biotin,
25 digoxigenin, His-Tag, Glutathion-S-Transferase, FLAG, GFP, myc-tag, and the like. In the case of a peptide or polypeptide, the tag is preferably at the N-terminus and/or C-terminus.

Suitable labels are any labels detectable by an appropriate detection method. Typical labels
30 include include gold particles, latex beads, acridan ester, luminol, ruthenium, enzymatically active labels, radioactive labels, magnetic labels ("e.g. magnetic beads", including paramagnetic and superparamagnetic labels), and fluorescent labels.

Enzymatically active labels include e.g. hydrogen peroxidase, alkaline phosphatase, and
35 derivatives thereof. Suitable substrates for detection of these enzymes include di-amino-benzidine (DAB), NBT-BCIP (4-nitro blue tetrazolium chloride and 5-bromo-4-chloro-3-indolyl-phosphate, available as ready-made stock solution from Roche Diagnostics), and

CDP-Star™ (Amersham). Suitable enzyme-substrate combinations result e.g. in colored reaction products, fluorescence or chemoluminescence. Typical fluorescent labels include Cy3, Cy5, Texas Red, Fluorescein, and the Alexa dyes (e.g. Alexa 568). Typical radioactive labels include ³⁵S, ¹²⁵I, ³²P and the like.

5

Suitable measurement methods according the present invention also include electrochemiluminescence (electro-generated chemiluminescence), ELISA (enzyme-linked immunosorbent assay), sandwich enzyme immune tests, turbidimetry, nephelometry, latex-enhanced turbidimetry or nephelometry, solid phase immune tests and 2D gel electrophoresis. If the position of the ADAM 12 protein (including its isoforms) in a 2D gel is known (e.g. identifiable by mass spectroscopy), then 2D gel electrophoresis may allow detection of ADAM 12 without use of a specifically binding ligand.

10

In another preferred embodiment, the peptide or polypeptide has a sequence which is selected from the sequence presented in SEQ ID NO: 8.

15

This embodiment relates to the isoform ADAM 12-S. ADAM 12-S is soluble and can be found at high concentrations in the blood. Therefore, ADAM 12-S can be measured easily using ligands binding to a peptide or polypeptide with a sequence according to SEQ ID NO: 4 and 8, particularly SEQ ID NO: 8. For specific detection of the short isoform (ADAM 12-S) or the long isoform (ADAM 12-L), binding of the ligand should be determined by regions unique to these isoforms, such as the C-terminal ends of ADAM 12-L (SEQ ID NO: 6) and ADAM 12-S (SEQ ID NO: 8). An example is an antibody raised against such a peptide or polypeptide. A suitable antibody has been described previously (Gilpin, B.J., Loechel, F., et al. (1998). A Novel, Secreted Form of Human ADAM 12 (Meltrin α) Provokes Myogenesis in Vivo. Journal of Biological Chemistry 273(1), 157-166). Said antibody is directed against a region (encoded by base pairs 2000-2433 according to Fig. 1 of Gilpin et al., 1998, cited above), which contains the first 4 amino acids (EARQ) of the C-terminal end unique to ADAM 12-S. However, in some samples, such as blood, ADAM 12-S is the most prevalent isoform and can be measured with a ligand directed against a region common to both isoforms.

20

25

30

In another preferred embodiment, the ligand is selected from the group consisting of

- a) KB-R7785 or a derivative thereof;

- b) TIMP-1, TIMP-2, TIMP-3, IGFBP-3, IGFBP-5, HB-EGF, α_2 -macroglobulin, PKC- δ , α -actinin-1, α -actinin-2, src, Grb-2, or syndecan-4;
- c) antibodies;
- d) nucleic acid or protein aptamers;
- 5 e) fragments or derivatives of any of the substances defined in b), c), or d).

Several ligands binding to ADAM 12 are known in the art, including small molecules (KB-R7785 (Asakura, M., Kitakaze, M., Takashima, S., Liao, Y. (2002). Cardiac hypertrophy is inhibited by antagonism of ADAM12 processing of HB-EGF: Metalloproteinase inhibitors as a new therapy. *Nature Medicine* 8: 35-40), polypeptides or proteins (TIMP-3, IGFBP-3, IGFBP-5, HB-EGF, α_2 -macroglobulin), and antibodies (see also Gilpin, B.J., et al. (1998), cited above). A polyclonal antibody against ADAM 12 is commercially available from Serotec (Cat. No. AHP758). Another antibody, directed against the pro-domain of ADAM-12, is available from Biomol (Cat. No. SA-378). The pro-domain is a domain that is cleaved off during activation of ADAM-12. Methods to generate antibodies or aptamers are well-known in the art. Identification and production of suitable antibodies or aptamers is also offered by commercial suppliers.

Notably, HB-EGF and IGFBPs, such as IGFBP-3 and IGFBP-5, are not only ligands of ADAM 12, but also substrates of the ADAM 12 proteolytic activity.

The person skilled in the art is familiar with methods to develop derivatives of the ligands according to a) to d) with higher affinity or specificity. For example, random mutations can be introduced into the proteins or substitutions into other molecules. These derivatives can then be tested for binding according to the screening procedure described above.

The term "antibody" as used herein includes both polyclonal and monoclonal antibodies, as well as fragments thereof, such as Fv, Fab and F(ab)₂ fragments that are capable of binding antigen or hapten. The present invention also includes "humanized" hybrid antibodies wherein amino acid sequences of a non-human donor antibody exhibiting a desired antigen-specificity are combined with sequences of a human acceptor antibody. The donor sequences will usually include at least the antigen-binding amino acid residues of the donor but may comprise other structurally and/or functionally relevant amino acid residues of the donor antibody as well. Such hybrids can be prepared by several methods well known in the art.

In a preferred embodiment, the ligand is directed against a fragment of any of the substances defined in b), c), or d). It is well-known that binding of antibodies, nucleic acids, peptides, or polypeptides occurs via specific binding regions, so-called binding sites, epitopes, or binding pockets. Such binding regions can be determined by methods known in the art, e.g. mutational analysis or NMR spectroscopy. Thus, fragments including only said binding regions can be determined. Those fragments may bind similarly or better than the original antibody, nucleic acid, peptide, polypeptide. Such fragments can also be used according to the present invention.

In a preferred embodiment, the above described use of nucleic acid molecules or ligands for diagnosis is done in conjunction with a diagnostic agent for the measurement of expression of any of the genes or proteins selected from the group consisting of (a) EPAS-1/HIF-2 α , (b) neurokinin B, (c) TIMP-1, (d) VEGFR-1, (e) VEGF, (f) IGFBP-1, (g) IGFBP-3, (h) matrix metalloproteinase-2, (i) leptin, (j) PAI-1, (k) IGF-1, (l) angiopoietin-2, (m) PIGF, (n) HLA-G, (o) HB-EGF, (p) TGF β -3 (q) decorin, and/or a diagnostic tool for the measurement of blood pressure or protein content of the urine. Comparative results for some of these genes are shown in Example 8.

Suitable diagnostic agents for the measurement of expression of said genes or proteins are known to the person skilled in the art. Suitable diagnostic agents can be defined analogously as above for ADAM 12. In particular, suitable diagnostic agents may comprise nucleic acids hybridizing to said genes or transcripts, antibodies, aptamers, and other ligands binding specifically to said proteins or the nucleic acids encoding said proteins.

Suitable antibodies are commercially available for most of the above defined proteins and can be found through public databases known to person skilled in the art, e.g. www.biocompare.com. Examples for suppliers are given in the following: EPAS-1/HIF-2 α (Abcam, Novus-Biologicals), neurokinin B (United States Biological), TIMP-1 (Affinity BioReagents (ABR)), VEGFR-1 (Chemicon, Lab Vision, Upstate, CalTag), VEGF (Anogen, Assay Designs Inc., BD Biosciences Pharmingen, Chemicon, Novus Biologicals, and others), IGFBP-1 (Cell Sciences, Novus Biologicals), IGFBP-3 (BD Biosciences Pharmingen, Novus Biologicals), matrix metalloproteinase-2 (Affinity BioReagents (ABR), Chemicon, Lab Vision), leptin (Affinity BioReagents (ABR), BD Pharmingen, BioSource International, Chemicon, Peprotech, and others), IGF-1 (Novus Biologicals), PAI-1 (BD Biosciences Pharmingen, United States Biological), angiopoietin-2 (Chemicon, Zymed), PIGF (Cell Sciences), HLA-G (Novus Biologicals, Serotec Inc.), HB-EGF (R&D Systems), TGF β -3 (Genex Bioscience), decorin (United States Biological).

Diagnostic tools for the measurement of blood pressure or protein content of the urine are known to the person skilled in the art. For example, the protein content of the urine may be measured by dip sticks.

5

In another preferred embodiment, a nucleic acid as defined above, a ligand as defined above, or, additionally, a diagnostic agent as defined above is present on an array.

10

Said array contains at least two nucleic acids as defined above, or ligands as defined above, or, additionally, diagnostic agents.

15

According to the present invention, the term "array" refers to a solid-phase or gel-like carrier upon which at least two compounds are attached or bound in one-, two- or three-dimensional arrangement. Such arrays are generally known to the person skilled in the art and typically generated on glass microscope slides, specially coated glass slides such as polycation-, nitrocellulose- or biotin-coated slides, cover slips, and membranes such as, for example, membranes based on nitrocellulose or nylon.

20

The aforementioned array may include a bound ligand, an above-defined nucleic acid, or at least two cells expressing at least one ligand or above-defined nucleic acid. In addition, said arrays may include a diagnostic agent or at least two cells expressing such diagnostic agent. Such arrays can be employed for diagnosis of preeclampsia and related syndromes.

25

The invention further relates to a method of producing arrays as defined above, wherein at least one above defined nucleic acid or ligand is bound to the carrier material in addition to other nucleic acids, ligands, or, additionally, diagnostic agents. Preferably, diagnostic agents for at least five, more preferably at least eight, additional marker genes or proteins are additionally bound to the carrier material.

30

Methods of producing such arrays, for example based on solid-phase chemistry and photolabile protective groups, are generally known (US 5,744,305). Such arrays can also be brought into contact with substances or substance libraries and tested for interaction, for example for binding or change of confirmation. Therefore, arrays comprising a peptide or polypeptide as defined above may be used for identifying ligands binding specifically to

35

said peptides or polypeptides.

The invention further relates to a method for preparing an array immobilized on a support material for diagnosis of preeclampsia or a related syndrome, in which at least two nucleic acids, at least two polypeptides or at least two antibodies or antibody fragments, and/or at least two cells or at least one of the aforementioned components in combination with other components relevant to preeclampsia and related disorders are used for preparation. The arrays produced by such a method can be employed for diagnosis of preeclampsia or a related syndrome.

In another preferred embodiment, diagnosis relates to the diagnosis of syndromes related to preeclampsia.

According to the present invention, syndromes "related" to preeclampsia include other pregnancy-associated disorders or syndromes with similar aetiology and/or symptoms as preeclampsia. Particularly included are eclampsia, intrauterine growth retardation, "missed abortion", superimposed gestosis, and HELLP syndrome.

In another embodiment, the present invention also relates to a method of diagnosis in which the above defined nucleic acid molecules, ligands, or, additionally, diagnostic agents or tools are used.

Using the above-defined nucleic acid molecules or the above-defined peptides or polypeptides, it is possible to identify suitable ligands by methods known in the art.

Therefore, the present invention also relates to the use of any of the nucleic acid molecules as defined above or the use of peptides or polypeptides comprising peptides or polypeptides as defined above for the identification of ligands binding specifically to said nucleic acids, peptides or polypeptides.

Suitable methods may comprise the steps of binding ADAM 12 polypeptides or suitable fragments thereof (such as the isoform-specific C-termini) or a candidate for a ligand to a solid phase, contacting the polypeptide with the ligand and detecting the binding of the ligand to the polypeptide. Examples comprise affinity chromatography, yeast-two-hybrid screens and enzyme-linked immuno-sorbent assays (ELISA). Identification of suitable ligands is also offered by commercial suppliers (e.g. Graffinity, Heidelberg), including the identification of binding antibodies, aptamers, and small molecules.

Therefore, the present invention also relates to a method for the identification of ligands binding specifically to any of the peptides or polypeptides as defined above, comprising the steps of contacting the polypeptide with at least one candidate for a ligand, and measuring the binding of the candidate for a ligand to the polypeptide. Said method may further comprise the steps of providing the polypeptide and providing at least one candidate for a ligand. Said peptides or polypeptides may be provided by organic synthesis or recombinant expression using the nucleic acid molecules as defined above. The peptides or polypeptides may also be isolated or purified from body fluids or tissue from the dead or alive human or animal body. Suitable purification methods are well known in the art and include salt fractionation, size exclusion chromatography, ion exchange chromatography, reverse phase chromatography, and affinity chromatography (with or without help of suitable tags, such as His-tags and the like). The isolated or purified peptides or polypeptides may contain modifications, such as glycosylation, against which a suitable ligand or candidate for a ligand may be directed.

After contacting the polypeptide with the candidate for a ligand, one or more optional steps of washing can be included.

The method can be carried out in the solid phase or in the liquid phase. If the method is carried out in the solid phase it may comprise a step of binding the polypeptide or a fragment thereof or the test substance to a solid phase. However, the step of contacting the polypeptide with at least one candidate for a ligand may still be carried out in a liquid, particularly in an aqueous solution. Examples for such methods are affinity chromatography and enzyme-linked immuno-sorbent assays (ELISA).

In another preferred embodiment, the method may be carried out in the liquid phase. Examples for such methods may include UV-crosslinking, immunoprecipitation, nuclear magnetic resonance spectroscopy (NMR, also available commercially, e.g. by Combinature Biopharm).

In another preferred embodiment, a method for identification of ligands may comprise the steps of (a) coupling a peptide or polypeptide comprising a peptide or polypeptide as defined above with a further detection moiety, (b) coupling the candidate for a ligand with a second detection moiety that is capable of interacting with the first detection moiety, (c) contacting the polypeptide of step (a) with the candidate for a ligand of step (b), preferably in a liquid, more preferably an aqueous solution, (d) measuring the interaction of the

detection moieties and thereby detecting the binding of the candidate for a ligand to the polypeptide.

An aqueous solution according to the present invention also relates to a cytosolic solution.
5 Examples for such methods include fluorescent resonance energy transfer (FRET) and yeast-two-hybrid screens. The interaction between the detection moieties may be either covalent or non-covalent, preferably non-covalent. Examples for suitable detection moieties include fluorescent molecules, such as green fluorescent protein (GFP) and its derivatives (Miyawaki, A., Tsien, R.Y., Monitoring protein conformations and interactions
10 by fluorescence resonance energy transfer between mutants of green fluorescent protein. Methods Enzymol. 2000; vol. 327, pp. 472-500) and transcription factors such as Gal4 and VP-16. Another optional step for a method of identification of ligands may comprise the step of comparing the binding of the candidate for a ligand to a related polypeptide, i.e. to another ADAM, such as ADAM 19, ADAM 33, or ADAM 2, to determine the specificity
15 of binding.

In another embodiment, the present invention relates to a method for diagnosis of preeclampsia or a related syndrome comprising the steps of (a) bringing a tissue sample or body fluid sample in contact with a nucleic acid as defined above or a specifically binding
20 ligand as defined above, and (b) detecting the binding of the nucleic acid or ligand. Preferably, the method is carried out in vitro.

Optionally, the method may include a step of obtaining a tissue sample or body fluid sample. Another optional step includes isolating nucleic acids or proteins contained in the
25 sample.

In another embodiment, the present invention relates to the use of a nucleic acid as defined above, a ligand as defined above, optionally in combination with a diagnostic agent or diagnostic tool as defined above, for the manufacture of a diagnostic for the diagnosis of
30 preeclampsia or a related syndrome.

In another embodiment, the present invention relates to a diagnostic containing a nucleic acid molecule, ligand, or, additionally, diagnostic agent as defined above. Such a diagnostic may be used for the diagnosis of preeclampsia or a related syndrome. The
35 nucleic acid molecule, ligand, or, additionally, diagnostic agent in the diagnostic may be labeled or unlabeled. The nucleic acid, ligand or diagnostic agent may also be present on a suitable carrier, e.g. a dipstick.

In another embodiment, the present invention relates to a diagnostic kit containing a nucleic acid molecule, ligand, or, additionally, diagnostic agent and/or diagnostic tool as defined above. Such a diagnostic kit may be used for the diagnosis of preeclampsia or a related syndrome.

The nucleic acid molecule, ligand, or, additionally, diagnostic agent in the kit may be labeled or unlabeled. Additionally, the kit may contain specifically binding secondary or higher order ligands (e.g. antibodies) and auxiliary reagents and tools as appropriate and known in the art. Furthermore, the kit may contain control reagents, such as agents for measuring total RNA, total protein content, or the expression level of "housekeeping" genes or proteins, such as EF-1 α , β -tubulin, cyclophilin B, HPRT, GAPDH, and others. The assays to be performed with the kit may be liquid phase assays as well as solid phase assays (i.e. with one or more reagents immobilized on a support). The kit may also comprise suitable tools for obtaining tissue or body fluid samples, particularly blood samples.

In another embodiment, the present invention also relates to a diagnostic or a diagnostic kit, wherein said nucleic acid, ligand or, additionally, diagnostic agent is present on an array. Preferably, diagnostic agents for at least five, more preferably at least eight, additional marker genes or proteins are additionally bound to the carrier material.

In another aspect, the invention relates to a method for the preparation of a diagnostic or a diagnostic kit, wherein a ligand of the peptides or polypeptides as defined above is identified according to the methods as defined above, synthesized in adequate amounts, and formulated into the diagnostic or diagnostic kit.

Similarly, said ligand may provided by any of the methods described above and formulated into the diagnostic or diagnostic kit.

The term "adequate amounts" of the ligand relates to any amounts sufficient for detection and/or measurement of said peptides or polypeptides. Suitable amounts depend on the particular method of detection or measurement, and are known to the person skilled in the art.

With respect to treatment, the problem according to the present invention is solved by use of a nucleic acid molecule comprising a nucleic acid selected from the group consisting of

- a) a nucleic acid with a sequence as presented in SEQ ID NO: 1 or 3;
- b) a fragment of any of the nucleic acids as defined in a);
- c) a nucleic acid with the antisense sequence of any of the sequences defined in a), or b);
- d) single-stranded or double-stranded RNA with a sequence corresponding to any of the sequences defined in a), b), or c);
- e) siRNA with a sequence defined in a), b), c), or d)

10 for the manufacture of a medicament for the treatment of preeclampsia or a related syndrome.

The problem is also solved by use of an inhibitor of the biological activity of a peptide or polypeptide with a sequence selected from the group consisting of

- a) an amino acid sequence as presented in SEQ ID No. 2 or 4;
- b) an amino acid sequence exhibiting a sequence identity with any of the sequences according to a) of at least 85% over 100 residues;
- c) a fragment of any of the sequences as defined above

20 for the manufacture of a medicament for the treatment of preeclampsia or a related syndrome.

25 These nucleic acid molecules, nucleic acids, peptides, polypeptides and sequences are in the following included in the group of "above defined" nucleic acid molecules, nucleic acids, peptides, polypeptides or sequences.

Preferably, the nucleic acid molecules, nucleic acids, peptides, or polypeptides have been purified and/or isolated.

30 As a consequence of the present invention, it is known that upregulation of ADAM 12, particularly ADAM 12-S, plays a significant role in preeclampsia and related syndromes. Therefore, treatment by inhibiting expression or biological activity of ADAM 12, particularly ADAM 12-S, is an object of the present invention.

35 A particular advantage of the present invention is that inhibition does not have to be complete. Already a reduction of the biological activity of ADAM 12, preferably to near

normal levels, can be sufficient for treatment. Thus, inhibition is also understood as a reduction of the biological activity of ADAM 12. Inhibition may also relate to modulation of ADAM 12 biological activity, such as modifying its substrate binding specificity or protease activity. It will be appreciated that inhibition of the biological activity can also be exerted through inhibition of expression, including inhibition of transcription, translation, or a decrease of transcript or protein stability.

According to the present invention, the term "inhibitor" relates to any substance inhibiting the biological activity of ADAM 12 or ADAM 12-S, particularly inhibiting the proteolytic or metalloproteolytic activity of ADAM 12 or ADAM 12-S. It also relates to inhibiting the binding of ADAM 12 or ADAM 12-S to endogenous substrates, such as HB-EGF or IGFBPs. It will be appreciated that all substrates of the proteolytic activity of ADAM 12 may be used as competitive inhibitors, including suitable peptides derived from those substrates. The person skilled in the art is familiar with identifying such peptides.

Preferably, the medicament is an inhibitor of expression or biological activity of ADAM 12 or ADAM 12-S. Inhibition of expression includes inhibition of transcription, translation, or a decrease of transcript or protein stability. Inhibition of biological activity particularly relates to inhibition of substrate binding activity or proteolytic activity.

A medicament according to the present invention preferably comprises additionally a suitable pharmaceutically acceptable carrier. The medicament may additionally comprise other carrier substances, preferably starch, lactose, fats, stearin acid, alcohol, physiological NaCl-solutions or further additives, in particular stabilizers, preservatives, dyes and flavourings. The medicament may also comprise other suitable substances. For example, an RNA or siRNA containing medicament may contain substances which stabilize RNA, particularly double-stranded RNA.

The above defined nucleic acid molecules are preferably used to inhibit expression of the corresponding endogenous genes. Nucleic acids which can be used to inhibit expression of the endogenous genes are known to the person skilled in the art. For example, peptide nucleic acids, comprising sequences as defined above, can suppress expression of the corresponding endogenous gene by forming DNA triplex structures with the gene. Other nucleic acids, such as antisense RNAs, siRNAs, morpholino oligonucleotides, or ribozymes, can be used to interfere with RNA transcribed from the endogenous gene. In

particular, small interfering RNAs (siRNA) can be used for inhibition of expression of the endogenous gene.

5 In a preferred embodiment, the nucleic acids have sequences shown in SEQ ID NO: 1 or 3, which correspond to those presented in GenBank Acc. No. AF023476 or AF023477.

10 In another preferred embodiment, fragments of any of the nucleic acids as defined above are used, preferably fragments with sequences located in SEQ ID NO: 5 or SEQ ID NO: 7. Suitable fragments are known to the person skilled in the art. For example, suitable fragments may be a peptide nucleic acid or oligonucleotides or nucleic acid fragments encoding siRNAs (also described in Tuschl, Nature Biotechnology, Vol. 20, pp. 446-44).

15 In another preferred embodiment, said nucleic acids have the antisense sequence of any of the sequences defined above. Such nucleic acids may comprise peptide nucleic acids (PNAs), antisense morpholino oligonucleotides, or antisense or double-stranded RNAs. Therefore, in another preferred embodiment, said nucleic acids may be single-stranded or double-stranded RNA with sequences corresponding to any of the sequences defined above.

20 In another preferred embodiment, the above-defined nucleic acid is an siRNA with sequences as defined above.

25 The inhibition of a specific target gene in mammals is achieved by the introduction of an siRNA-molecule having a sequence that is specific for the target gene into the mammalian cell. The siRNAs comprise a first and a second RNA strand, both hybridized to each other, wherein the sequence of the first RNA strand is an oligonucleotide corresponding to one of the sequences as defined above for therapy and wherein the sequence of the second RNA strand is the antisense-strand of the first RNA strand. The siRNA-molecules may possess a characteristic 2- or 3-nucleotide 3'-overhang. Each strand of the siRNA molecule
30 preferably has a length of 19 to 31 nucleotides.

The siRNAs can be introduced into the mammalian cell by any suitable known method of cell transfection, particularly lipofection, electroporation, or microinjection, be it in vitro or in vivo.

The RNA oligonucleotides can be generated and hybridized to each other in vitro or in vivo according to any of the known RNA synthesis methods, in order to obtain siRNA.

5 In another embodiment, the invention relates to the use of a nucleic acid molecule, wherein the nucleic acid molecule is contained in at least one nucleic acid expression vector which is capable of producing a double-stranded RNA-molecule comprising a sense-RNA-strand and an antisense-RNA-strand under suitable conditions, wherein each RNA-strand, independently from the other, has a length of 19 to 31 nucleotides.

10 In this alternative method (also described in Tuschl, Nature Biotechnology, Vol. 20, pp. 446-448), vector systems capable of producing siRNAs instead of the siRNAs themselves are introduced into the mammalian cell for downregulating gene expression.

15 In a preferred embodiment, the first and the second RNA strand of the siRNA may have, independently from the other, a length of 19 to 25 nucleotides, more preferred of 20 to 25 nucleotides, and most preferred of 20 to 22 nucleotides.

20 In another preferred embodiment, the first and the second RNA strand of the siRNA may have, independently from the other, a length of 26 to 30 nucleotides, more preferred of 26 to 28 nucleotides, and most preferred of 27 nucleotides.

In another embodiment, the present invention relates to a medicament comprising an above-defined nucleic acid molecule.

25 Administration of the medicament can be carried out by known methods. For therapeutic applications, the medicament may be in form of a solution, in particular an injectable solution, a cream, ointment, tablet, suspension, granulate or the like. The medicament may be administered in any suitable way, in particular by injection, by oral, nasal, rectal application. The medicament may particularly be administered parenteral, that means
30 without entering the digestion apparatus, for example by subcutaneous injection. In particular, the medicament may also be injected intravenously in the form of solutions for infusions or injections. Other suitable administration forms may be direct administrations on the skin in the form of creams, ointments, sprays and other transdermal therapeutic

substances or in the form of inhalative substances, such as nose sprays, aerosoles or in the form of microcapsules or implantates.

Preferably, the inhibitor is administered in pharmaceutically effective amount. As used
5 herein, a "pharmaceutically effective amount" of an inhibitor is an amount effective to achieve the desired physiological result, either in cells treated *in vitro* or in a subject treated *in vivo*. Specifically, a pharmaceutically effective amount is an amount sufficient to positively influence, for some period of time, one or more clinically defined pathological effects or symptoms associated with preeclampsia. The pharmaceutically effective amount
10 may vary depending on the specific inhibitor selected, and is also dependent on a variety of factors and conditions related to the subject to be treated and the severity of the disorder or symptom. For example, if the inhibitor is to be administered *in vivo*, factors such as age, weight, gestational age, and general health of the patient as well as dose response curves and toxicity data obtained in pre-clinical animal tests would be among the factors to be
15 considered. If the inhibitor is to be contacted with cells *in vitro*, one would also design a variety of pre-clinical *in vitro* studies to assess parameters like uptake, half-life, dose, toxicity etc. The determination of a pharmaceutically effective amount for a given agent is well within the ability of those skilled in the art. Preferably, the inhibitor is present in a concentration of 0,1 to 50% per weight of the pharmaceutical composition, more
20 preferably 10 to 30%.

In another embodiment, the present invention relates to a method of treatment of preeclampsia or a related syndrome comprising the use of the above-identified nucleic acid molecules.

25

Treatment of preeclampsia or a related syndrome relates to treatment of the disorder or syndrome itself as well as treatment of symptoms associated with said disease or syndrome. Such symptoms may include intravascular coagulation, blood platelet destruction, high blood pressure, edema, proteinuria, or placental abruption. Treatment
30 according to the present invention also relates to prophylaxis.

Treatment according to the present invention may be accompanied by treatment approaches known in the art, for example blood pressure control (particularly by

administration of alpha-receptor blockers (e.g. hydralazine) or ebrantil), tocolysis (particularly for increase of placental blood flow), administration of magnesium, and administration of cortisone.

- 5 In another preferred embodiment, the inhibitor is a disintegrin domain metalloproteinase inhibitor, particularly KB-R7785, a TIMP, particularly TIMP-3 or a fragment thereof, α_2 -macroglobulin, or an antibody directed against ADAM 12.

10 In a preferred embodiment, such an inhibiting antibody is directed against the proteolytic domain of ADAM 12, particularly the zink binding domain, more particularly including amino acid positions 350-361.

Preferably, the inhibitor should be specific for ADAM 12, ADAM 12-L, or ADAM 12-S. The criteria for specificity as defined above for ligands apply analogously.

15

In a preferred embodiment, the peptide or polypeptide is a fragment with a sequence consisting of or located in SEQ ID NO: 8. According to this embodiment, the inhibitor is directed specifically against a peptide or polypeptide consisting of or located in SEQ ID NO: 8. An example is an antibody raised against such a peptide or polypeptide. It is known
20 that such antibodies, though in this case binding only to the C-terminus of ADAM 12-S, can sterically inhibit the biological activity of the entire ADAM 12-S protein. The same is true for aptamers and other inhibitors. Suitable inhibitors can be found by identifying ligands binding specifically to ADAM 12-S and subsequently screening them as candidates for inhibitors. Analogously, inhibitors directed against the C-terminus of ADAM 12-L
25 (SEQ ID NO: 6) may be identified.

In another preferred embodiment, the above-defined nucleic acid or inhibitor is used in conjunction with HB-EGF. Preferably, HB-EGF has been purified or isolated. Preferably, HB-EGF is formulated into the medicament defined above.

30

In another preferred embodiment, the medicament is for treatment of symptoms of preeclampsia or a related syndrome, particularly for treatment or prevention of intravascular coagulation, blood platelet destruction, high blood pressure, edema, glomerular endotheliosis, proteinuria, or placental abruption.

35

Symptoms of preeclampsia and related syndromes include any undesired effects the disorders may have on the mother or fetus. Particularly, symptoms relate to symptoms

caused by a high level of ADAM 12 in the blood, particularly the maternal blood. More particularly, such symptoms include intravascular coagulation (thrombosis), blood platelet destruction, or high blood pressure (particularly above 140/90 mm Hg), edema (swelling of face and/or hands), glomerular endotheliosis, or proteinuria (particularly more than 0.3 g/24 hours or a level of 2+ by dipstick test).

Treatment may also be restricted to the mother's blood. For example, IgM antibodies or other molecules of high molecular weight have only a limited capability of passing the placental blood barrier. Therefore, in a preferred embodiment, the inhibitor may be an IgM or any other molecule not capable of passing the placental blood barrier, particularly being of similar or higher molecular weight as an IgM. For example, an identified inhibitor, such as an aptamer, may be coupled covalently or non-covalently to a high molecular weight residue, such as a polypeptide or dextran.

In another aspect, the present invention relates to the use of the nucleic acids, peptides or polypeptides as defined above for identification of inhibitors of the biological activity of the above-defined nucleic acids, peptides or polypeptides, particularly for identification of inhibitors of the proteolytic activity or substrate-binding activity of such peptides or polypeptides.

The person skilled in the art is familiar with such use of nucleic acids, peptides or polypeptides. Examples for suitable methods are described in the following.

Therefore, in another aspect, the present invention relates to a method for identification of inhibitors of any of the peptides or polypeptides as defined above, comprising the steps of (a) contacting said peptide or polypeptide with a suitable substrate, e.g. HB-EGF, (b) measuring the decrease in processing of the substrate in the presence as compared to the absence of a candidate for an inhibitor molecule.

Suitable substrates include all substrates of biological, preferably enzymatic, more preferably proteolytic activity. Suitable substrates include HB-EGF and IGFBPs, particularly IGFBP-3 and IGFBP-5. Processing of the substrate relates preferably to enzymatic modification, more preferably proteolytic modification or cleavage of the substrate. Measurement of processing can be done e.g. by measuring the decrease in substrate (educt) and the increase in modified substrate (product). Measurement of processing can involve indirect measurement, e.g. via ligands, labeled or unlabeled (see above) binding to the substrate or modified substrate.

Candidates for inhibitor molecules can be any nucleic acid, peptide, polypeptide or other substance. In particular, candidates may include peptides derived from TIMP, IGFBP-3, IGFBP-5, HB-EGF, α_2 -macroglobulin, antibodies, nucleic acid or protein aptamers or fragments or derivatives of any of these substances. Other candidates for inhibitor molecules may comprise derivatives of KB-R7785. The person skilled in the art is familiar with methods of generating such derivatives, either by methods of combinatorial chemistry or introducing mutations into the respective nucleic acids or polypeptides.

In a preferred embodiment, the present invention relates to the candidate for inhibitor being a substrate or ligand as defined above. Any substrate of ADAM 12 or ADAM 12-S or ligand as defined above is also considered a candidate for an inhibitor.

In another aspect, the present invention relates to a method for the preparation of a pharmaceutical composition, wherein an inhibitor of the nucleic acids, peptides or polypeptides as defined above is identified according to the method as defined above, synthesized in adequate amounts, and formulated into a pharmaceutical composition.

Suitable methods to synthesize the inhibitor molecules are known in the art. For example, peptides or polypeptides can be synthesized by recombinant expression (see also above), and antibodies can be obtained from hybridoma cell lines or immunized animals. Small molecules can be synthesized according to any known organic synthesis method.

Similarly, said inhibitor may be provided by any of the screening methods described above and formulated into a pharmaceutical composition.

The term "adequate amounts" of the inhibitor relates to any amounts sufficient for inhibiting expression or biological activity of ADAM-12 in the human or animal body, preferably the human body. In particular, adequate amounts relate to pharmaceutically effective amounts.

Description of the Figures

Fig. 1 Increased ADAM 12-S expression in preeclamptic human placenta. A, gestational age (GA, weeks + days) of the study subjects (Pat. No.). B, Semi-quantitative RT-PCR analysis for the levels of ADAM 12-S transcripts in human placenta using cyclophilin B for normalization. Note the increased level of ADAM 12-S transcript in samples of preeclamptic patients (Preecl.) as compared samples of healthy

mothers (Ctrl.). The increase is visible both in the fetal (f) and maternal (m) part of the placenta. Patient 22 was pregnant with twins, and samples 22-1 and 22-2 were derived from the placentae belonging to each of the twins.

5 Fig. 2 Increased ADAM 12-S expression in preeclamptic human placenta. Shown are relative mRNA levels for ADAM 12-S in the fetal (f) or maternal (m) part of healthy (Ctrl.) and preeclamptic (Preecl.) human placentas as determined by real time PCR using elongation factor EF-1 α for normalization. Note the increased level of mRNA in samples of preeclamptic patients as compared samples of healthy pregnant women. Y-axis: level of fluorescence of the PCR product (which is correlated with the level of mRNA in the sample), X-axis: Sample numbers, corresponding to the patient numbers (Pat No.) in Fig. 1A. Patient 22 was pregnant with twins, and samples 22-1 and 22-2 were derived from each placenta of twin siblings.

15 The invention is further illustrated by the following examples, which are not intended to limit the scope of the invention in any way.

EXAMPLE 1

20 Placentae from different gestational age from preeclamptic patients as well as from healthy control patients were dissected and collected within less than 15 min after delivery. Preeclampsia was diagnosed according to the generally accepted criteria (blood pressure above 140/90 mm Hg and proteinuria of more than 0.3 g/24 hours or a level of 2+ by dipstick test). 10-20 mm thick sections were dissected and the fetal and maternal side were collected separately. Tissue was either immediately snap frozen or fixed in 4% paraformaldehyde for 12 hours. The fixed tissue was washed twice in phosphate-buffered saline (PBS, composition of 1l: 8g NaCl (137 mM), 0,2g KCl (2,7 mM), 1.44g Na₂HPO₄·2H₂O (8,1 mM), 0,24g KH₂PO₄ (1,76mM), pH 7.4), transferred to 70% ethanol and paraffin embedded. For RNA preparation the tissue was incubated for 24 hrs in 30 RNALater (Ambion) at 4°C and then frozen at -80°C.

EXAMPLE 2

Total RNA was isolated from frozen placental tissue, either from healthy or preeclamptic pregnant women of different gestational age, with Trizol (Invitrogen, Karlsruhe, Germany) according to the manufacturer's recommendations. The integrity and purity of the total RNA was checked on a 1% agarose gel with denaturing loading buffer or on the RNA 6000 Nano Chip (Agilent Technology, Germany). Total RNA quality and concentration

was determined by absorption spectroscopy between 220 and 320 nm ($1 \text{ OD}^{260} = 40 \mu\text{g/ml}$ ssRNA).

EXAMPLE 3

5 The microarrays contained 1600 nonredundant, sequence-validated human cDNAs (RZPD set, Berlin, (Fritz, B., et al. (2002). Microarray-based copy number and expression profiling in dedifferentiated and pleomorphic liposarcoma. *Cancer Res*, vol. 62: 2993-8). The cDNAs had been selected on their putative involvement in tumorigenesis and were PCR amplified as described previously (Wrobel, G., et al. (2003). Optimization of high-
10 density cDNA-microarray protocols by 'design of experiments'. *Nucleic Acids Res*, vol. 31: e67, and Schlingemann, J., et al. (2003). Profile of gene expression induced by the tumor promotor TPA in murine epithelial cells. *Int J Cancer* 104(6), p. 699-708), purified by isopropanol precipitation and dissolved in 20 μl FBNC (Formamid, betaine and nitrocellulose) spotting buffer (Wrobel, G., et al. (2003), see above). The resulting cDNA
15 fragments had an average length between 500-1500 bp.

EXAMPLE 4

DNA spotting was performed on Poly-Prep poly-L-lysine coated substrates (Sigma Chemical Co.; However, also other substrates known in the art may be used, such as QF
20 Epoxy substrates (Quantifoil Microtools, Jena, Germany)) using an OmniGrid Microarrayer (GeneMachines, San Carlos, CA) equipped with Stealth SMP3 Micro Spotting Pins (Telechem). Spot centres were 250 μm apart. The arrays were provided with positive control spots of the Spot Report System (Stratagene, Amsterdam, The Netherlands) as well as a variety of additional negative controls, e.g., mouse Cot-1 DNA
25 (Invitrogen), yeast tRNA (Sigma Chemical Co.) and poly-dA (Amersham Biosciences). Furthermore, Cy3- and Cy5-labeled DNA was included to yield specific fluorescent landmarks on the arrays. DNA immobilisation to the glass was achieved by 3.5 hours of incubation at 80°C, followed by 2x irradiation at 254 nm with an energy of 120 mJ/cm^2 in a Stratalinker Model 2400 UV illuminator (Stratagene).

EXAMPLE 5

30-40 μg of total RNA (as obtained according to Example 2) was used for microarray hybridisation. Reverse transcription was performed in accordance to Wrobel et al., 2003 using an Omniscript RT Kit (QIAGEN) and C3/Cy5-dUTPs (NEN Life Science Products,
35 Köln, Germany). Subsequently, Cy3- and Cy5-labeled cDNA samples were pooled (age matched placental tissue from healthy and preeclamptic placental tissues), purified and

concentrated to 10 µl using a Microcon YM-30 PCR filter unit (Millipore, Eschborn, Germany), as previously described.

EXAMPLE 6

- 5 31 µl of DIG Easy Hybridisation buffer (Roche Diagnostics, Mannheim, Germany) were added to each cDNA pool and incubated for 15-60 min at room temperature. Denaturation was carried out at 65°C for 3 min. Hybridisation was performed for 16h at 37°C in ArrayIt Hybridisation cassettes (TeleChem International, Sunnyvale) in a waterbath. Thereafter the slides were washed at room temperature with (i) 1x SSC, 0.1% (w/v) SDS for 10 min for 3
10 times; (ii) 0.1x SSC, 0.1% SDS for 10 min; (iii) H₂O for 3 min; (iv) 70% ethanol for 3 min; (v) 90% ethanol for 3 min; (vi) 100% ethanol for 3 min and (vii) finally dried with compressed air.

EXAMPLE 7

- 15 Microarrays were scanned and analysed using a GenePix 4000A microarray scanner (Axon Instruments, Inc., Union City, NJ), controlled by the software GenePix Pro 3.0 (Axon Instruments, Inc.). Individual data reports for all conducted experiments (including, e.g., median fluorescence intensities of all spots and local background) were exported and further analysed using the R software environment for statistical computing (<http://www.r-project.org>) as well as Microsoft Excel 2000. Here, data sets for spots not recognized by
20 the GenePix analysis Software were excluded from further considerations. Additionally, all remaining data sets were ranked according to spot homogeneity (as assayed by the ratio of median and mean fluorescence intensities), spot intensity and the standard deviation of log ratios for replicate spots. Those data points ranked among the lower 50%, and thus
25 possessing the lower reliability based on the criteria just described, were eliminated. Data sets that could not be verified, either in a colour switch experiment (reversed assignment of fluorophores) or on a different experimental day, were omitted as well. For each hybridisation, fluorescence ratios (Cy5/Cy3) were normalised by the median log ratio of all genes found in this experiment. Accurate differential expression values for each gene were
30 obtained by calculating the average of the normalised ratios of all independent hybridisations, successfully identifying that gene, and reversing the ratios from colour switch experiments prior to this operation.

EXAMPLE 8

- 35 Comparative microarray analyses were performed for with the microarray as described in Example 3. The following pairs of samples with the indicated gestational ages (gestational weeks + days) were used:

Pair No.	healthy (control)	preeclamptic
1	41+3	41+0
2	38+2	38+2
3	34+0	35+1 (1)
4	34+0	35+1 (2)
5	34+0	33+2

- 5 35+1 (1) and 35+1 (2) designate samples derived from each placenta of twin siblings (patient 22, see legend to Fig. 1 and 2).

It was found that ADAM-12 was the most strongly regulated gene. Shown below are the data of Pair No. 2. Expression levels were calculated from two independent experiments.

10

Part of placenta	induced (upregulated)			repressed (downregulated)		
	fetal	maternal	both	fetal	maternal	both
No. of genes induced more than 2-fold	35	27	27	35	35	34
ADAM-12	38-fold	44.5-fold		-	-	
TIMP-1	8.5-fold	12-fold		-	-	
TIMP-2	13-fold	11.5-fold		-	-	
EPAS/HIF-2 α	4-fold	2.5-fold		-	-	
IGFBP-1	-	-		less than 2-fold	7.5-fold	
IGFBP-3	-	-		2.6-fold	5.7-fold	
TEK	-	-		4.8-fold	2.8-fold	
decorin	-	-		less than 2-fold	6.5-fold	

ADAM 6 and ADAM 9 were also included on the microarray, but were not found to be induced or repressed by more than 2-fold.

EXAMPLE 9

For RT-PCR analysis 2-5 µg total RNA from healthy and preeclamptic placental tissue of different gestational age pretreated with RQ1-DNase (Promega, Madison, WI) were reverse transcribed in a reaction volume of 20 µl using AMV-RT (Promega) and an oligo-d(T) (Promega) according to the manufacturer's recommendations, 2-3 µl of the RT reaction together with primers depicted in the following table were used for PCR detection.

Primer Pairs used for RT-PCR analysis		
Gene Name	Forward primer 5'	Reverse primer 3'
ADAM 12-L	CCTGTGACCTCCCAGAGTTCTGC (SEQ ID NO: 9)	GAATTCCTATGGTTAAACCTTGG (SEQ ID NO: 10)
ADAM 12-L	CTTGCTGCCGATTGTGGTTTAT (SEQ ID NO: 11)	AGTTGACTGGGGCTGAGGGACATT (SEQ ID NO: 12)
ADAM 12-S	CCTGTGACCTCCCAGAGTTCTGC (SEQ ID NO: 9)	GGACTCTGCAGCTTCCTGCCTTG (SEQ ID NO: 14)
ADAM 12-S	TCGGGGCTCAGGAGGGGAAGT (SEQ ID NO: 15)	GGCTGCCAAGGCGGAGGTGTC (SEQ ID NO: 16)
human cyclophilinB	GGCCGGGTGATCTTTGGTCTCTTC (SEQ ID NO: 17)	CCCGGCTGTCTGTCTTGGTGCTCT (SEQ ID NO: 18)
human EF-1α	CTGCTGAGATGGGAAAGGGCT (SEQ ID NO: 19)	TTCAGGATAATCACCTGAGCA (SEQ ID NO: 20)

Human cyclophilin B served as an internal control for the quality and quantity of the cDNA. Amplified probes were separated on 1.5% agarose gels in the presence of ethidium bromide and visualised by using the Eagle Eye system (Stratagene). To obtain a calibration graph, cDNA PCR product from human elongation factor EF-1α was serially diluted in H₂O (five 10-fold dilutions from 0.01 pg to 10 pg). To standardize the amount of sample cDNA, an endogenous control amplicon was used (EF-1α).

EXAMPLE 10

Semi-quantitative RT-PCR was performed on placental tissue samples obtained from 4 healthy and 3 preeclamptic patients (Fig. 1A). Typical results of the RT-PCR

amplifications are shown in Fig. 1B. Samples 22-1 and 22-2 were obtained from the same patient.

A comparison of data from healthy and preeclamptic women is shown in the graph of Fig. 2. Overall, the RT-PCR experiments demonstrate a 4 to 5-fold increase in ADAM 12-S transcript in placental tissue of both maternal and fetal origin.

EXAMPLE 11

The following procedure was used to obtain RNA from plasma samples. Peripheral blood samples were collected with informed consent of the patients. 5 to 10 ml blood samples were collected in EDTA-containing tubes and centrifuged at 4000 rpm for 15 min at 4°C. Plasma was collected in new tubes. The plasma samples were recentrifuged at 13000 rpm for 15 min at 4°C.

For analysis of RNA derived from plasma, the supernatants obtained in the above procedure are transferred to fresh tubes and can be processed according to a previously reported protocol, which is herewith incorporated by reference (Ng, E.K.O., Leung, T.N., et al. (2003). The concentration of circulating Corticotropin-releasing hormone RNA in maternal plasma is increased in preeclampsia. Clin. Chem. 49, 727-731, see particularly p. 728-729 and Ng, E.K.O., Tsui, N.B.Y., et al. (2003). Messenger RNA of placental origin is readily detectable in maternal plasma. Proc Natl Acad Sci USA, vol. 100(8):4748-53, see particularly p. 4748-4749).

EXAMPLE 12

For Real-time Quantitative PCR (RQ PCR), the QuantiTect™ SYBR Green PCR Kit (Qiagen, Hilden, Germany) was used to amplify ADAM-12S specific templates from cDNAs of placental RNA of different patients according to the manufacturer's instructions. The amount of the amplified templates in each sample was analysed with the MJ Research Opticon™ (Biozym, Oldendorf, Germany) calculating against a standard PCR with elongation factor EF-1α, amplified from different dilutions.

Quantitative assessment of the DNA amplification was detected via the dye SYBR Green, the fluorescence of which is strongly enhanced upon the binding to the dsDNA. The RQ-PCR reactions were carried out in 50 µl using 1 µl of the RT reaction at a primer concentration of 0.3 µM, respectively, according to the manufacturer's recommendations for QuantiTect™ SYBR Green PCR Kit reagents (Qiagen, Hilden, Germany). The oligonucleotides used for real time PCR are listed in example 9. For thermal cycling the

following conditions were applied: one step at 94°C 15 min, then 40 cycles of 30 sec at 94°C, 1 min at 56°C and 1:30 min at 72°C and a final step at 72°C for 3 min. Melting curve was achieved from 50 up to 93°C within reading every 0.5°C hold for 10 sec between reads to obtain melting curves of the final RQ-PCR products. This is necessary because SYBR Green fluorescence may also be derived from side products such as primer dimers.

To obtain a calibration graph, cDNA PCR product from elongation factor EF-1 α was serially diluted in H₂O (five 10-fold dilutions from 0.01 pg to 10 pg). To standardize the amount of sample cDNA, an endogenous control amplicon was used (EF-1 α).

21. Okt. 2003

- 1 -

Deutsches Krebsforschungszentrum
Universität HeidelbergOctober 17, 2003
DK62208EP HN/FLZ/ROE/hen/hmü**Patent Claims**

5

1. Use of a nucleic acid molecule comprising a nucleic acid selected from the group consisting of

10

- a) a nucleic acid with a sequence as presented in SEQ ID NO: 1 or 3;
- b) a nucleic acid with a sequence that exhibits a sequence identity with any of the sequences defined in a) of at least 70% over 300 residues;
- c) a nucleic acid which is capable of hybridizing with the nucleic acid as defined in a), or b) under conditions of medium or high stringency;
- d) a nucleic acid with the antisense-sequence of any of the sequences defined
- 15 in a), b) or c);
- e) a fragment of any of the nucleic acids as defined in a), b), c), or d);
- f) an RNA corresponding to any of the sequences defined in a), b), c), d) or e);

15

for diagnosis of preeclampsia or a related syndrome.

20

2. The use according to claim 1, wherein the fragment has a sequence consisting of or being part of SEQ ID NO: 7, 13, 14, 15, or 16, or wherein the fragment has a sequence consisting of or being part of SEQ ID NO: 5, 10, 11, or 12.

25

3. Use of a ligand binding specifically to a peptide or polypeptide with a sequence selected from the group consisting of

30

- a) an amino acid sequence as presented in SEQ ID NO: 2 or 4;
- b) an amino acid sequence exhibiting a sequence identity with any of the amino acid sequences according to a) of at least 85% over 100 amino acid residues;
- c) a fragment of any of the sequences defined above

for diagnosis of preeclampsia or a related syndrome.

35

4. The use according to claim 3, wherein the sequence is selected from SEQ ID NO: 8.

5. The use according to any of claims 3 to 4, wherein the ligand is selected from the group consisting of

- a) KB-R7785 or a derivative thereof;
- b) TIMP-1, TIMP-2, TIMP-3, IGFBP-3, IGFBP-5, HB-EGF, α_2 -macroglobulin, PKC- δ , α -actinin, α -actinin-2, src, Grb-2, or syndecan-4
- c) antibodies;
- d) nucleic acid or protein aptamers;
- e) fragments or derivatives of any of the substances defined in b), c), or d).

6. The use according to any of claims 1 to 5, wherein the nucleic acid or ligand is used in conjunction with a diagnostic agent for the measurement of expression of any of the genes or proteins selected from the group consisting of

- a) EPAS-1/HIF-2 α ,
- b) neurokinin B
- c) TIMP-1,
- d) VEGFR-1,
- e) VEGF,
- f) IGFBP-1
- g) IGFBP-3,
- h) matrix metalloproteinase-2,
- i) leptin,
- j) PAI-1,
- k) IGF-1,
- l) angiopoietin-2,
- m) decorin,
- n) PIGF,
- o) HLA-G
- p) HB-EGF
- q) TGF- β 3

and/or diagnostic tools for the measurement of blood pressure or protein content of the urine.

7. The use according to any of claims 1 to 6, wherein the nucleic acid, ligand or, additionally, diagnostic agent is present on an array.
8. Use of the nucleic acid molecule as defined in any of claims 1 to 2, or of a peptide or polypeptide comprising a peptide or polypeptide as defined in any of claims 3 or 4 for the identification of ligands binding specifically to said nucleic acid, peptide or polypeptide.
9. A method for the identification of ligands binding specifically to a peptide or polypeptide as defined in in any of claims 3 to 4, comprising the following steps:
 - a) contacting the polypeptide with at least one candidate for a ligand;
 - b) measuring the binding of the candidate for a ligand to the polypeptide.
10. A method for diagnosis of preeclampsia or a related syndrome comprising the following steps:
 - a) bringing a biopsy or bodily fluid sample into contact with a nucleic acid as defined in any of claims 1 to 2 or a specifically binding ligand as defined in claims 3 to 5, and
 - b) detecting the binding of the nucleic acid or ligand.
11. Use of a nucleic acid as defined in any of claims 1 to 2, or a ligand as defined in any of claims 3 to 5, optionally in combination with a diagnostic agent or diagnostic tool as defined in claim 6, for the manufacture of a diagnostic for the diagnosis of preeclampsia or a related syndrome.
12. A diagnostic containing a nucleic acid or ligand, or, additionally, diagnostic agent or tool as defined in any of claims 1 to 5.
13. A diagnostic kit containing a nucleic acids or ligand as defined in any of claims 1 to 5.
14. The diagnostic or diagnostic kit according to any of claims 12 to 13, wherein the nucleic acid, ligand or, additionally, diagnostic agent is present on an array.

15. Use of a nucleic acid molecule comprising a nucleic acid selected from the group consisting of

- a) a nucleic acid with a sequence as presented in SEQ ID NO: 1 or 3;
- b) a fragment of any of the nucleic acids as defined in a);
- c) a nucleic acid with the antisense sequence of any of the sequences defined in a), or b);
- d) single-stranded or double-stranded RNA, preferably siRNA, with a sequence corresponding to any of the sequences defined in a), b), or c);

for the manufacture of a medicament for the treatment of preeclampsia or a related syndrome.

16. The use according to claim 15, wherein the fragment consists of or is located in SEQ ID NO: 5 or 7.

17. Use of an inhibitor of the biological activity of a peptide or polypeptide with a sequence selected from the group consisting of

- a) an amino acid sequence as presented in SEQ ID No. 2 or 4;
- b) an amino acid sequence exhibiting a sequence identity with any of the sequences according to a) of at least 85% over 100 residues;
- c) a fragment of any of the sequences as defined above

for the manufacture of a medicament for the treatment of preeclampsia or a related syndrome.

18. The use according to claim 17, wherein the fragment has a sequence consisting of or being part of SEQ ID NO: 8.

19. The use according to any of claims 17 to 18, wherein the inhibitor is a disintegrin domain metalloproteinase inhibitor, particularly KB-R7785, a TIMP, particularly TIMP-3 or a fragment thereof, α_2 -Macroglobulin, or an antibody directed against ADAM 12.

20. The use according to any of claims 17 to 19, wherein additionally HB-EGF is used for the manufacture of the medicament.
- 5 21. The use according to any of claims 15 to 20, wherein the medicament is for treatment of symptoms of preeclampsia or a related syndrome, particularly for treatment or prevention of intravascular coagulation, blood platelet destruction, placental abruption, or high blood pressure.
- 10 22. Use of a nucleic acid, peptide, or polypeptide as defined in any of claims 1 to 4 for identification of an inhibitor of said nucleic acid, peptide or polypeptide, particularly for identification of an inhibitor of the proteolytic activity or substrate-binding activity of said peptide or polypeptide.
- 15 23. A method for identification of an inhibitor of the biological activity of a peptide or polypeptide as defined in any of claims 17 to 18, comprising the following steps:
- a) contacting said peptide or polypeptide with a suitable substrate, e.g. HB-EGF,
 - b) measuring the decrease in processing of the substrate in the presence as compared to the absence of a candidate for an inhibitor molecule
- 20 24. The method according to claim 22, wherein the candidate for an inhibitor is a substrate or a ligand of a peptide or polypeptide as defined in any of claims 3 to 4.
- 25 25. A method for the preparation of a pharmaceutical composition, wherein an inhibitor of the nucleic acids, peptides, or polypeptides as defined in any of claims 17 to 18 is identified according to the methods as defined in claim 22 or 23, synthesized in adequate amounts, and formulated into a pharmaceutical composition.

21. Okt. 2003

- 1 -

Deutsches Krebsforschungszentrum
Universität Heidelberg

October 17, 2003
DK62208EP HN/FLZ/ROE/hen

Summary

5

The present invention relates to the significant role of ADAM 12, ADAM 12-L, and ADAM 12-S in preeclampsia and related disorders. The invention further relates to the use of said genes and gene products, ligands binding to those, and modulators of activity of the gene products in diagnosis and treatment of preeclampsia and related syndromes.

21. Okt. 2003

Fig. 1

A

Pat. No.	GA	Ctrl.	Preecl.
5	38+2	+	-
6	37+2	+	-
20	34	+	-
21	41+3	+	-
14	38+2	-	+
16	33+2	-	+
22	35+1	-	+

B





Fig. 2



SEQUENCE LISTING

EPO - Munich
79

21. Okt. 2003

<110> Deutsches Krebsforschungszentrum

<120> Use of ADAM 12 for diagnosis and therapy of preeclampsia

<130> DK62208EP

<160> 20

<170> PatentIn version 3.1

<210> 1

<211> 5062

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (307)..(3036)

<223>

```

<400> 1
cactaacgct cttcctagtc cccgggccaa ctcggacagt ttgctcattt attgcaacgg      60
tcaaggctgg cttgtgccag aacggcgcg cgcgcgacgca cgcacacaca cgggggggaaa      120
cttttttaaa aatgaaaggc tagaagagct cagcggcggc gcgggccgtg cgcgagggct      180
ccggagctga ctcgccgagg caggaaatcc ctccggtcgc gacgcccggc cccgctcggc      240
gcccgcgtgg gatggtgcag cgctcgccgc cggggccgag agctgctgca ctgaaggccg      300
gcgacg atg gca gcg cgc ccg ctg ccc gtg tcc ccc gcc cgc gcc ctc      348
      Met Ala Ala Arg Pro Leu Pro Val Ser Pro Ala Arg Ala Leu
      1          5          10
ctg ctc gcc ctg gcc ggt gct ctg ctc gcg ccc tgc gag gcc cga ggg      396
Leu Leu Ala Leu Ala Gly Ala Leu Leu Ala Pro Cys Glu Ala Arg Gly
15          20          25          30
gtg agc tta tgg aac caa gga aga gct gat gaa gtt gtc agt gcc tct      444
Val Ser Leu Trp Asn Gln Gly Arg Ala Asp Glu Val Val Ser Ala Ser
      35          40          45
gtt cgg agt ggg gac ctc tgg atc cca gtg aag agc ttc gac tcc aag      492
Val Arg Ser Gly Asp Leu Trp Ile Pro Val Lys Ser Phe Asp Ser Lys

```

50	55	60	
aat cat cca gaa gtg ctg aat att cga cta caa cgg gaa agc aaa gaa Asn His Pro Glu Val Leu Asn Ile Arg Leu Gln Arg Glu Ser Lys Glu 65 70 75			540
ctg atc ata aat ctg gaa aga aat gaa ggt ctc att gcc agc agt ttc Leu Ile Ile Asn Leu Glu Arg Asn Glu Gly Leu Ile Ala Ser Ser Phe 80 85 90			588
acg gaa acc cac tat ctg caa gac ggt act gat gtc tcc ctc gct cga Thr Glu Thr His Tyr Leu Gln Asp Gly Thr Asp Val Ser Leu Ala Arg 95 100 105 110			636
aat tac acg gta att ctg ggt cac tgt tac tac cat gga cat gta cgg Asn Tyr Thr Val Ile Leu Gly His Cys Tyr Tyr His Gly His Val Arg 115 120 125			684
gga tat tct gat tca gca gtc agt ctc agc acg tgt tct ggt ctc agg Gly Tyr Ser Asp Ser Ala Val Ser Leu Ser Thr Cys Ser Gly Leu Arg 130 135 140			732
gga ctt att gtg ttt gaa aat gaa agc tat gtc tta gaa cca atg aaa Gly Leu Ile Val Phe Glu Asn Glu Ser Tyr Val Leu Glu Pro Met Lys 145 150 155			780
agt gca acc aac aga tac aaa ctc ttc cca gcg aag aag ctg aaa agc Ser Ala Thr Asn Arg Tyr Lys Leu Phe Pro Ala Lys Lys Leu Lys Ser 160 165 170			828
gtc cgg gga tca tgt gga tca cat cac aac aca cca aac ctc gct gca Val Arg Gly Ser Cys Gly Ser His His Asn Thr Pro Asn Leu Ala Ala 175 180 185 190			876
aag aat gtg ttt cca cca ccc tct cag aca tgg gca aga agg cat aaa Lys Asn Val Phe Pro Pro Pro Ser Gln Thr Trp Ala Arg Arg His Lys 195 200 205			924
aga gag acc ctc aag gca act aag tat gtg gag ctg gtg atc gtg gca Arg Glu Thr Leu Lys Ala Thr Lys Tyr Val Glu Leu Val Ile Val Ala 210 215 220			972
gac aac cga gag ttt cag agg caa gga aaa gat ctg gaa aaa gtt aag Asp Asn Arg Glu Phe Gln Arg Glu Lys Asp Leu Glu Lys Val Lys 225 230 235			1020
cag cga tta ata gag att gct aat cac gtt gac aag ttt tac aga cca Gln Arg Leu Ile Glu Ile Ala Asn His Val Asp Lys Phe Tyr Arg Pro 240 245 250			1068
ctg aac att cgg atc gtg ttg gta ggc gtg gaa gtg tgg aat gac atg Leu Asn Ile Arg Ile Val Leu Val Gly Val Glu Val Trp Asn Asp Met 255 260 265 270			1116
gac aaa tgc tct gta agt cag gac cca ttc acc agc ctc cat gaa ttt Asp Lys Cys Ser Val Ser Gln Asp Pro Phe Thr Ser Leu His Glu Phe 275 280 285			1164
ctg gac tgg agg aag atg aag ctt cta cct cgc aaa tcc cat gac aat Leu Asp Trp Arg Lys Met Lys Leu Leu Pro Arg Lys Ser His Asp Asn 290 295 300			1212
gcg cag ctt gtc agt ggg gtt tat ttc caa ggg acc acc atc ggc atg Ala Gln Leu Val Ser Gly Val Tyr Phe Gln Gly Thr Thr Ile Gly Met 305 310 315			1260
gcc cca atc atg agc atg tgc acg gca gac cag tct ggg gga att gtc Ala Pro Ile Met Ser Met Cys Thr Ala Asp Gln Ser Gly Gly Ile Val 320 325 330			1308

atg Met 335	gac Asp	cat His	tca Ser	gac Asp	aat Asn 340	ccc Pro	ctt Leu	ggt Gly	gca Ala	gcc Ala 345	gtg Val	acc Thr	ctg Leu	gca Ala	cat His 350	1356
gag Glu	ctg Leu	ggc Gly	cac His	aat Asn 355	ttc Phe	ggg Gly	atg Met	aat Asn 360	cat His	gac Asp	aca Thr	ctg Leu	gac Asp	agg Arg 365	ggc Gly	1404
tgt Cys	agc Ser	tgt Cys	caa Gln 370	atg Met	gcg Ala	gtt Val	gag Glu	aaa Lys 375	gga Gly	ggc Gly	tgc Cys	atc Ile	atg Met 380	aac Asn	gct Ala	1452
tcc Ser	acc Thr	ggg Gly 385	tac Tyr	cca Pro	ttt Phe	ccc Pro	atg Met 390	gtg Val	ttc Phe	agc Ser	agt Ser	tgc Cys 395	agc Ser	agg Arg	aag Lys	1500
gac Asp 400	ttg Leu	gag Glu	acc Thr	agc Ser	ctg Leu	gag Glu 405	aaa Lys	gga Gly	atg Met	ggg Gly	gtg Val 410	tgc Cys	ctg Leu	ttt Phe	aac Asn	1548
ctg Leu 415	ccg Pro	gaa Glu	gtc Val	agg Arg	gag Glu 420	tct Ser	ttc Phe	ggg Gly	ggc Gly	cag Gln 425	aag Lys	tgt Cys	ggg Gly	aac Asn	aga Arg 430	1596
ttt Phe	gtg Val	gaa Glu	gaa Glu	gga Gly 435	gag Glu	gag Glu	tgt Cys	gac Asp	tgt Cys 440	ggg Gly	gag Glu	cca Pro	gag Glu	gaa Glu 445	tgt Cys	1644
atg Met	aat Asn	cgc Arg	tgc Cys 450	tgc Cys	aat Asn	gcc Ala	acc Thr	acc Thr 455	tgt Cys	acc Thr	ctg Leu	aag Lys	ccg Pro 460	gac Asp	gct Ala	1692
gtg Val	tgc Cys	gca Ala 465	cat His	ggg Gly	ctg Leu	tgc Cys	tgt Cys 470	gaa Glu	gac Asp	tgc Cys	cag Gln 475	ctg Leu 475	aag Lys	cct Pro	gca Ala	1740
gga Gly 480	aca Thr	gcg Ala	tgc Cys	agg Arg	gac Asp	tcc Ser 485	agc Ser	aac Asn	tcc Ser	tgt Cys	gac Asp 490	ctc Leu	cca Pro	gag Glu	ttc Phe	1788
tgc Cys 495	aca Thr	ggg Gly	gcc Ala	agc Ser	cct Pro 500	cac His	tgc Cys	cca Pro	gcc Ala 505	aac Asn 505	gtg Val	tac Tyr	ctg Leu	cac His	gat Asp 510	1836
ggg Gly	cac His	tca Ser	tgt Cys	cag Gln 515	gat Asp	gtg Val	gac Asp	ggc Gly	tac Tyr 520	tgc Cys	tac Tyr	aat Asn	ggc Gly	atc Ile 525	tgc Cys	1884
cag Gln	act Thr	cac His	gag Glu 530	cag Gln	cag Gln	tgt Cys	gtc Val	aca Thr 535	ctc Leu	tgg Trp	gga Gly	cca Pro	ggt Gly 540	gct Ala	aaa Lys	1932
cct Pro	gcc Ala	cct Pro 545	ggg Gly	atc Ile	tgc Cys	ttt Phe	gag Glu 550	aga Arg	gtc Val	aat Asn	tct Ser	gca Ala 555	ggt Gly	gat Asp	cct Pro	1980
tat Tyr 560	ggc Gly	aac Asn	tgt Cys	ggc Gly	aaa Lys	gtc Val 565	tcg Ser	aag Lys	agt Ser	tcc Ser	ttt Phe 570	gcc Ala	aaa Lys	tgc Cys	gag Glu	2028
atg Met 575	aga Arg	gat Asp	gct Ala	aaa Lys	tgt Cys 580	gga Gly	aaa Lys	atc Ile	cag Gln	tgt Cys 585	caa Gln	gga Gly	ggt Gly	gcc Ala	agc Ser 590	2076
cgg Arg	cca Pro	gtc Val	att Ile	ggt Gly 595	acc Thr	aat Asn	gcc Ala	gtt Val	tcc Ser 600	ata Ile	gaa Glu	aca Thr	aac Asn	atc Ile 605	ccc Pro	2124

ctg cag caa gga ggc cgg att ctg tgc cgg ggg acc cac gtg tac ttg Leu Gln Gln Gly Gly Arg Ile Leu Cys Arg Gly Thr His Val Tyr Leu 610 615 620	2172
ggc gat gac atg ccg gac cca ggg ctt gtg ctt gca ggc aca aag tgt Gly Asp Asp Met Pro Asp Pro Gly Leu Val Leu Ala Gly Thr Lys Cys 625 630 635	2220
gca gat gga aaa atc tgc ctg aat cgt caa tgt caa aat att agt gtc Ala Asp Gly Lys Ile Cys Leu Asn Arg Gln Cys Gln Asn Ile Ser Val 640 645 650	2268
ttt ggg gtt cac gag tgt gca atg cag tgc cac ggc aga ggg gtg tgc Phe Gly Val His Glu Cys Ala Met Gln Cys His Gly Arg Gly Val Cys 655 660 665 670	2316
aac aac agg aag aac tgc cac tgc gag gcc cac tgg gca cct ccc ttc Asn Asn Arg Lys Asn Cys His Cys Glu Ala His Trp Ala Pro Pro Phe 675 680 685	2364
tgt gac aag ttt ggc ttt gga gga agc aca gac agc ggc ccc atc cgg Cys Asp Lys Phe Gly Phe Gly Gly Ser Thr Asp Ser Gly Pro Ile Arg 690 700	2412
caa gca gat aac caa ggt tta acc ata gga att ctg gtg acc atc ctg Gln Ala Asp Asn Gln Gly Leu Thr Ile Gly Ile Leu Val Thr Ile Leu 705 710 715	2460
tgt ctt ctt gct gcc gga ttt gtg gtt tat ctc aaa agg aag acc ttg Cys Leu Leu Ala Ala Gly Phe Val Val Tyr Leu Lys Arg Lys Thr Leu 720 725 730	2508
ata cga ctg ctg ttt aca aat aag aag acc acc att gaa aaa cta agg Ile Arg Leu Leu Phe Thr Asn Lys Lys Thr Thr Ile Glu Lys Leu Arg 735 740 745 750	2556
tgt gtg cgc cct tcc cgg cca ccc cgt ggc ttc caa ccc tgt cag gct Cys Val Arg Pro Ser Arg Pro Pro Arg Gly Phe Gln Pro Cys Gln Ala 755 760 765	2604
cac ctc ggc cac ctt gga aaa ggc ctg atg agg aag ccg cca gat tcc His Leu Gly His Leu Gly Lys Gly Leu Met Arg Lys Pro Pro Asp Ser 770 775 780	2652
tac cca ccg aag gac aat ccc agg aga ttg ctg cag tgt cag aat gtt Tyr Pro Pro Lys Asp Asn Pro Arg Arg Leu Leu Gln Cys Gln Asn Val 785 790 795	2700
gac atc agc aga ccc ctc aac ggc ctg aat gtc cct cag ccc cag tca Asp Ile Ser Arg Pro Leu Asn Gly Leu Asn Val Pro Gln Pro Gln Ser 800 805 810	2748
act cag cga gtg ctt cct ccc ctc cac cgg gcc cca cgt gca cct agc Thr Gln Arg Val Leu Pro Pro Leu His Arg Ala Pro Arg Ala Pro Ser 815 820 825 830	2796
gtc cct gcc aga ccc ctg cca gcc aag cct gca ctt agg cag gcc cag Val Pro Ala Arg Pro Leu Pro Ala Lys Pro Ala Leu Arg Gln Ala Gln 835 840 845	2844
ggg acc tgt aag cca aac ccc cct cag aag cct ctg cct gca gat cct Gly Thr Cys Lys Pro Asn Pro Pro Gln Lys Pro Leu Pro Ala Asp Pro 850 855 860	2892
ctg gcc aga aca act cgg ctc act cat gcc ttg gcc agg acc cca gga Leu Ala Arg Thr Thr Arg Leu Thr His Ala Leu Ala Arg Thr Pro Gly 865 870 875	2940
caa tgg gag act ggg ctc cgc ctg gca ccc ctc aga cct gct cca caa	2988

Gln	Trp	Glu	Thr	Gly	Leu	Arg	Leu	Ala	Pro	Leu	Arg	Pro	Ala	Pro	Gln	
880						885					890					
tat	cca	cac	caa	gtg	ccc	aga	tcc	acc	cac	acc	gcc	tat	att	aag	tga	3036
Tyr	Pro	His	Gln	Val	Pro	Arg	Ser	Thr	His	Thr	Ala	Tyr	Ile	Lys		
895					900					905						
gaagccgaca	ccttttttca	acagtgaaga	cagaagtttg	cactatcttt	cagctccagt											3096
tggagttttt	tgtaccaact	tttaggattt	tttttaatgt	ttaaaacatc	attactataa											3156
gaactttgag	ctactgccgt	cagtgcgtgtg	ctgtgctatg	gtgctctgtc	tacttgcaca											3216
ggtacttgta	aattattaat	ttatgcagaa	tggtgattac	agtgcagtgc	gctgtagtag											3276
gcatttttac	catcactgag	ttttccatgg	caggaaggct	tggtgtgctt	ttagtatttt											3336
agtgaacttg	aaatatcctg	cttgatggga	ttctggacag	gatgtgtttg	ctttctgatc											3396
aaggccttat	tggaagcag	tcccccaact	acccccagct	gtgcttatgg	taccagatgc											3456
agctcaagag	atcccaagta	gaatctcagt	tgattttctg	gattcccat	ctcaggccag											3516
agccaagggg	cttcaggtcc	aggctgtgtt	tggctttcag	ggaggccctg	tgcccttgga											3576
caactggcag	gcaggctccc	agggacacct	gggagaaatc	tggcttctgg	ccaggaagct											3636
ttggtgagaa	cctgggttgc	agacaggaat	cttaaggtgt	agccacacca	ggatagagac											3696
tggaacacta	gacaagccag	aacttgaccc	tgagctgacc	agccgtgagc	atgtttggaa											3756
ggggtctgta	gtgtcactca	aggcgggtgct	tgatagaaat	gccaagcact	tctttttctc											3816
gctgtccttt	ctagagcact	gccaccagta	ggttatttag	cttgggaaag	gtggtgtttc											3876
tgtaagaaac	ctactgccc	ggcactgcaa	accgccacct	ccctatactg	cttggagctg											3936
agcaaatacac	cacaaactgt	aatacaatga	tcctgtattc	agacagatga	ggactttcca											3996
tgggaccaca	actattttca	gatgtgaacc	attaaccaga	tctagtcaat	caagtctgtt											4056
tactgcaagg	ttcaacttat	taacaattag	gcagactctt	tatgcttgca	aaaactacaa											4116
ccaatggaat	gtgatgttca	tgggtatagt	tcagtgtctgc	tatcattatt	cgtagatatt											4176
ggacaaagaa	ccttctctat	ggggcatcct	ctttttccaa	cttggctgca	ggaatcttta											4236
aaagatgctt	ttacagagt	ctgaacctat	ttcttaaaca	cttgcaacct	acctgttgag											4296
catcacagaa	tgtgataagg	aaatcaactt	gcttatcaac	ttcctaaata	ttatgagatg											4356
tggcttgggc	agcatcccct	tgaactcttc	actcttcaaa	tgcttgacta	gggagccatg											4416
tttcacaagg	tctttaaagt	gactaatggc	atgagaaata	caaaaatact	cagataaggt											4476
aaaatgccat	gatgcctctg	tcttctggac	tggttttcac	attagaagac	aattgacaac											4536
agttacataa	ttcactctga	gtgttttatg	agaaagcctt	cttttggggt	caacagtttt											4596
cctatgcttt	gaaacagaaa	aatatgtacc	aagaatcttg	gtttgccttc	cagaaaacaa											4656
aactgcattt	cactttcccg	gtgttcccca	ctgtatctag	gcaacatagt	attcatgact											4716
atggataaac	taaacacgtg	acacaacac	acacaaaagg	gaaccagct	ctaatacatt											4776
ccaactcgta	tagcatgcat	ctgtttattc	tatagttatt	aagttcttta	aaatgtaaag											4836
ccatgctgga	aaataatact	gctgagatac	atacagaatt	actgtaactg	attacacttg											4896

gtaattgtac taaagccaaa catatatata ctattaaaaa ggtttacaga attttatggt 4956
gcattacgtg ggcattgtct ttttagatgc ccaaattcctt agatctggca tgtttagccct 5016
tcctccaatt ataagaggat atgaacccaa aaaaaaaaaa aaaaaa 5062

<210> 2

<211> 909

<212> PRT

<213> Homo sapiens

<400> 2

Met Ala Ala Arg Pro Leu Pro Val Ser Pro Ala Arg Ala Leu Leu Leu
1 5 10 15

Ala Leu Ala Gly Ala Leu Leu Ala Pro Cys Glu Ala Arg Gly Val Ser
20 25 30

Leu Trp Asn Gln Gly Arg Ala Asp Glu Val Val Ser Ala Ser Val Arg
35 40 45

Ser Gly Asp Leu Trp Ile Pro Val Lys Ser Phe Asp Ser Lys Asn His
50 55 60

Pro Glu Val Leu Asn Ile Arg Leu Gln Arg Glu Ser Lys Glu Leu Ile
65 70 75 80

Ile Asn Leu Glu Arg Asn Glu Gly Leu Ile Ala Ser Ser Phe Thr Glu
85 90 95

Thr His Tyr Leu Gln Asp Gly Thr Asp Val Ser Leu Ala Arg Asn Tyr
100 105 110

Thr Val Ile Leu Gly His Cys Tyr Tyr His Gly His Val Arg Gly Tyr
115 120 125

Ser Asp Ser Ala Val Ser Leu Ser Thr Cys Ser Gly Leu Arg Gly Leu
130 135 140

Ile Val Phe Glu Asn Glu Ser Tyr Val Leu Glu Pro Met Lys Ser Ala
145 150 155 160

Thr Asn Arg Tyr Lys Leu Phe Pro Ala Lys Lys Leu Lys Ser Val Arg
165 170 175

Gly Ser Cys Gly Ser His His Asn Thr Pro Asn Leu Ala Ala Lys Asn
180 185 190

Val Phe Pro Pro Pro Ser Gln Thr Trp Ala Arg Arg His Lys Arg Glu
195 200 205

Thr Leu Lys Ala Thr Lys Tyr Val Glu Leu Val Ile Val Ala Asp Asn
 210 215 220
 Arg Glu Phe Gln Arg Gln Gly Lys Asp Leu Glu Lys Val Lys Gln Arg
 225 230 240
 Leu Ile Glu Ile Ala Asn His Val Asp Lys Phe Tyr Arg Pro Leu Asn
 245 250 255
 Ile Arg Ile Val Leu Val Gly Val Glu Val Trp Asn Asp Met Asp Lys
 260 265 270
 Cys Ser Val Ser Gln Asp Pro Phe Thr Ser Leu His Glu Phe Leu Asp
 275 280 285
 Trp Arg Lys Met Lys Leu Leu Pro Arg Lys Ser His Asp Asn Ala Gln
 290 295 300
 Leu Val Ser Gly Val Tyr Phe Gln Gly Thr Thr Ile Gly Met Ala Pro
 305 310 315 320
 Ile Met Ser Met Cys Thr Ala Asp Gln Ser Gly Gly Ile Val Met Asp
 325 330 335
 His Ser Asp Asn Pro Leu Gly Ala Ala Val Thr Leu Ala His Glu Leu
 340 345 350
 Gly His Asn Phe Gly Met Asn His Asp Thr Leu Asp Arg Gly Cys Ser
 355 360 365
 Cys Gln Met Ala Val Glu Lys Gly Gly Cys Ile Met Asn Ala Ser Thr
 370 375 380
 Gly Tyr Pro Phe Pro Met Val Phe Ser Ser Cys Ser Arg Lys Asp Leu
 385 390 395 400
 Glu Thr Ser Leu Glu Lys Gly Met Gly Val Cys Leu Phe Asn Leu Pro
 405 410 415
 Glu Val Arg Glu Ser Phe Gly Gly Gln Lys Cys Gly Asn Arg Phe Val
 420 425 430
 Glu Glu Gly Glu Glu Cys Asp Cys Gly Glu Pro Glu Glu Cys Met Asn
 435 440 445
 Arg Cys Cys Asn Ala Thr Thr Cys Thr Leu Lys Pro Asp Ala Val Cys
 450 455 460
 Ala His Gly Leu Cys Cys Glu Asp Cys Gln Leu Lys Pro Ala Gly Thr
 465 470 475 480

Ala Cys Arg Asp Ser Ser Asn Ser Cys Asp Leu Pro Glu Phe Cys Thr
 485 490 495
 Gly Ala Ser Pro His Cys Pro Ala Asn Val Tyr Leu His Asp Gly His
 500 505 510
 Ser Cys Gln Asp Val Asp Gly Tyr Cys Tyr Asn Gly Ile Cys Gln Thr
 515 520 525
 His Glu Gln Gln Cys Val Thr Leu Trp Gly Pro Gly Ala Lys Pro Ala
 530 535 540
 Pro Gly Ile Cys Phe Glu Arg Val Asn Ser Ala Gly Asp Pro Tyr Gly
 545 550 555 560
 Asn Cys Gly Lys Val Ser Lys Ser Ser Phe Ala Lys Cys Glu Met Arg
 565 570 575
 Asp Ala Lys Cys Gly Lys Ile Gln Cys Gln Gly Gly Ala Ser Arg Pro
 580 585 590
 Val Ile Gly Thr Asn Ala Val Ser Ile Glu Thr Asn Ile Pro Leu Gln
 595 600 605
 Gln Gly Gly Arg Ile Leu Cys Arg Gly Thr His Val Tyr Leu Gly Asp
 610 615 620
 Asp Met Pro Asp Pro Gly Leu Val Leu Ala Gly Thr Lys Cys Ala Asp
 625 630 635 640
 Gly Lys Ile Cys Leu Asn Arg Gln Cys Gln Asn Ile Ser Val Phe Gly
 645 650 655
 Val His Glu Cys Ala Met Gln Cys His Gly Arg Gly Val Cys Asn Asn
 660 665 670
 Arg Lys Asn Cys His Cys Glu Ala His Trp Ala Pro Pro Phe Cys Asp
 675 680 685
 Lys Phe Gly Phe Gly Gly Ser Thr Asp Ser Gly Pro Ile Arg Gln Ala
 690 695 700
 Asp Asn Gln Gly Leu Thr Ile Gly Ile Leu Val Thr Ile Leu Cys Leu
 705 710 715 720
 Leu Ala Ala Gly Phe Val Val Tyr Leu Lys Arg Lys Thr Leu Ile Arg
 725 730 735
 Leu Leu Phe Thr Asn Lys Lys Thr Thr Ile Glu Lys Leu Arg Cys Val
 740 745 750

Arg Pro Ser Arg Pro Pro Arg Gly Phe Gln Pro Cys Gln Ala His Leu
 755 760 765
 Gly His Leu Gly Lys Gly Leu Met Arg Lys Pro Pro Asp Ser Tyr Pro
 770 775 780
 Pro Lys Asp Asn Pro Arg Arg Leu Leu Gln Cys Gln Asn Val Asp Ile
 785 790 795 800
 Ser Arg Pro Leu Asn Gly Leu Asn Val Pro Gln Pro Gln Ser Thr Gln
 805 810 815
 Arg Val Leu Pro Pro Leu His Arg Ala Pro Arg Ala Pro Ser Val Pro
 820 825 830
 Ala Arg Pro Leu Pro Ala Lys Pro Ala Leu Arg Gln Ala Gln Gly Thr
 835 840 845
 Cys Lys Pro Asn Pro Pro Gln Lys Pro Leu Pro Ala Asp Pro Leu Ala
 850 855 860
 Arg Thr Thr Arg Leu Thr His Ala Leu Ala Arg Thr Pro Gly Gln Trp
 865 870 875 880
 Glu Thr Gly Leu Arg Leu Ala Pro Leu Arg Pro Ala Pro Gln Tyr Pro
 885 890 895
 His Gln Val Pro Arg Ser Thr His Thr Ala Tyr Ile Lys
 900 905

<210> 3
 <211> 3328
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (307)..(2523)
 <223>

<400> 3
 cactaacgct cttcctagtc cccgggccaa ctcggacagt ttgctcattt attgcaacgg 60
 tcaaggctgg cttgtgccag aacggcgcg cgcgcgacgca cgcacacaca cggggggaaa 120
 cttttttaaa aatgaaaggc tagaagagct cagcggcggc gcgggccgtg cgcgagggct 180
 ccggagctga ctcgccgagg caggaaatcc ctccggtcgc gacgcccggc cccgctcggc 240

gcccgcgtgg gatggtgcag cgctcgccgc cgggcccag agctgctgca ctgaaggccg	300
gcgacg atg gca gcg cgc ccg ctg ccc gtg tcc ccc gcc cgc gcc ctc Met Ala Ala Arg Pro Leu Pro Val Ser Pro Ala Arg Ala Leu 1 5 10	348
ctg ctc gcc ctg gcc ggt gct ctg ctc gcg ccc tgc gag gcc cga ggg Leu Leu Ala Leu Ala Gly Ala Leu Leu Ala Pro Cys Glu Ala Arg Gly 15 20 25 30	396
gtg agc tta tgg aac gaa gga aga gct gat gaa gtt gtc agt gcc tct Val Ser Leu Trp Asn Glu Gly Arg Ala Asp Glu Val Val Ser Ala Ser 35 40 45	444
gtt cgg agt ggg gac ctc tgg atc cca gtg aag agc ttc gac tcc aag Val Arg Ser Gly Asp Leu Trp Ile Pro Val Lys Ser Phe Asp Ser Lys 50 55 60	492
aat cat cca gaa gtg ctg aat att cga cta caa cgg gaa agc aaa gaa Asn His Pro Glu Val Leu Asn Ile Arg Leu Gln Arg Glu Ser Lys Glu 65 70 75	540
ctg atc ata aat ctg gaa aga aat gaa ggt ctc att gcc agc agt ttc Leu Ile Ile Asn Leu Glu Arg Asn Glu Gly Leu Ile Ala Ser Ser Phe 80 85 90	588
acg gaa acc cac tat ctg caa gac ggt act gat gtc tcc ctc gct cga Thr Glu Thr His Tyr Leu Gln Asp Gly Thr Asp Val Ser Leu Ala Arg 95 100 105 110	636
aat tac acg gta att ctg ggt cac tgt tac tac cat gga cat gta cgg Asn Tyr Thr Val Ile Leu Gly His Cys Tyr Tyr His Gly His Val Arg 115 120 125	684
gga tat tct gat tca gca gtc agt ctc agc acg tgt tct ggt ctc agg Gly Tyr Ser Asp Ser Ala Val Ser Leu Ser Thr Cys Ser Gly Leu Arg 130 135 140	732
gga ctt att gtg ttt gaa aat gaa agc tat gtc tta gaa cca atg aaa Gly Leu Ile Val Phe Glu Asn Glu Ser Tyr Val Leu Glu Pro Met Lys 145 150 155	780
agt gca acc aac aga tac aaa ctc ttc cca gcg aag aag ctg aaa agc Ser Ala Thr Asn Arg Tyr Lys Leu Phe Pro Ala Lys Lys Leu Lys Ser 160 165 170	828
gtc cgg gga tca tgt gga tca cat cac aac aca cca aac ctc gct gca Val Arg Gly Ser Cys Gly Ser His His Asn Thr Pro Asn Leu Ala Ala 175 180 185 190	876
aag aat gtg ttt cca cca ccc tct cag aca tgg gca aga agg cat aaa Lys Asn Val Phe Pro Pro Pro Ser Gln Thr Trp Ala Arg Arg His Lys 195 200 205	924
aga gag acc ctc aag gca act aag tat gtg gag ctg gtg atc gtg gca Arg Glu Thr Leu Lys Ala Thr Lys Tyr Val Glu Leu Val Ile Val Ala 210 215 220	972
gac aac cga gag ttt cag agg caa gga aaa gat ctg gaa aaa gtt aag Asp Asn Arg Glu Phe Gln Arg Gln Gly Lys Asp Leu Glu Lys Val Lys 225 230 235	1020
cag cga tta ata gag att gct aat cac gtt gac aag ttt tac aga cca Gln Arg Leu Ile Glu Ile Ala Asn His Val Asp Lys Phe Tyr Arg Pro 240 245 250	1068
ctg aac att cgg atc gtg ttg gta ggc gtg gaa gtg tgg aat gac atg Leu Asn Ile Arg Ile Val Leu Val Gly Val Glu Val Trp Asn Asp Met 255 260 265 270	1116

gac Asp	aaa Lys	tgc Cys	tct Ser	gta Val 275	agt Ser	cag Gln	gac Asp	cca Pro	ttc Phe 280	acc Thr	agc Ser	ctc Leu	cat His	gaa Glu 285	ttt Phe	1164
ctg Leu	gac Asp	tgg Trp	agg Arg 290	aag Lys	atg Met	aag Lys	ctt Leu	cta Leu 295	cct Pro	cgc Arg	aaa Lys	tcc Ser	cat His 300	gac Asp	aat Asn	1212
gcg Ala	cag Gln	ctt Leu 305	gtc Val	agt Ser	ggg Gly	gtt Val	tat Tyr 310	ttc Phe	caa Gln	ggg Gly	acc Thr	acc Thr 315	atc Ile	ggc Gly	atg Met	1260
gcc Ala	cca Pro 320	atc Ile	atg Met	agc Ser	atg Met	tgc Cys 325	acg Thr	gca Ala	gac Asp	cag Gln	tct Ser 330	ggg Gly	gga Gly	att Ile	gtc Val	1308
atg Met 335	gac Asp	cat His	tca Ser	gac Asp	aat Asn 340	ccc Pro	ctt Leu	ggt Gly	gca Ala	gcc Ala 345	gtg Val	acc Thr	ctg Leu	gca Ala	cat His 350	1356
gag Glu	ctg Leu	ggc Gly	cac His	aat Asn 355	ttc Phe	ggg Gly	atg Met	aat Asn	cat His 360	gac Asp	aca Thr	ctg Leu	gac Asp	agg Arg 365	ggc Gly	1404
tgt Cys	agc Ser	tgt Cys	caa Gln 370	atg Met	gcg Ala	gtt Val	gag Glu	aaa Lys 375	gga Gly	ggc Gly	tgc Cys	atc Ile	atg Met 380	aac Asn	gct Ala	1452
tcc Ser	acc Thr	ggg Gly 385	tac Tyr	cca Pro	ttt Phe	ccc Pro	atg Met 390	gtg Val	ttc Phe	agc Ser	agt Ser	tgc Cys 395	agc Ser	agg Arg	aag Lys	1500
gac Asp	ttg Leu 400	gag Glu	acc Thr	agc Ser	ctg Leu	gag Glu 405	aaa Lys	gga Gly	atg Met	ggg Gly	gtg Val 410	tgc Cys	ctg Leu	ttt Phe	aac Asn	1548
ctg Leu 415	ccg Pro	gaa Glu	gtc Val	agg Arg	gag Glu 420	tct Ser	ttc Phe	ggg Gly	ggc Gly	cag Gln 425	aag Lys	tgt Cys	ggg Gly	aac Asn	aga Arg 430	1596
ttt Phe	gtg Val	gaa Glu	gaa Glu	gga Gly 435	gag Glu	gag Glu	tgt Cys	gac Asp	tgt Cys 440	ggg Gly	gag Glu	cca Pro	gag Glu	gaa Glu 445	tgt Cys	1644
atg Met	aat Asn	cgc Arg	tgc Cys 450	tgc Cys	aat Asn	gcc Ala	acc Thr	acc Thr 455	tgt Cys	acc Thr	ctg Leu	aag Lys	ccg Pro 460	gac Asp	gct Ala	1692
gtg Val	tgc Cys	gca Ala 465	cat His	ggg Gly	ctg Leu	tgc Cys	tgt Cys 470	gaa Glu	gac Asp	tgc Cys	cag Gln	ctg Leu 475	aag Lys	cct Pro	gca Ala	1740
gga Gly 480	aca Thr	gcg Ala	tgc Cys	agg Arg	gac Asp	tcc Ser 485	agc Ser	aac Asn	tcc Ser	tgt Cys	gac Asp 490	ctc Leu	cca Pro	gag Glu	ttc Phe	1788
tgc Cys 495	aca Thr	ggg Gly	gcc Ala	agc Ser	cct Pro 500	cac His	tgc Cys	cca Pro	gcc Ala	aac Asn 505	gtg Val	tac Tyr	ctg Leu	cac His	gat Asp 510	1836
ggg Gly	cac His	tca Ser	tgt Cys	cag Gln 515	gat Asp	gtg Val	gac Asp	ggc Gly	tac Tyr 520	tgc Cys	tac Tyr	aat Asn	ggc Gly	atc Ile 525	tgc Cys	1884
cag Gln	act Thr	cac His	gag Glu 530	cag Gln	cag Gln	tgt Cys	gtc Val	aca Thr 535	ctc Leu	tgg Trp	gga Gly	cca Pro	ggt Gly 540	gct Ala	aaa Lys	1932

cct gcc cct ggg atc tgc ttt gag aga gtc aat tct gca ggt gat cct Pro Ala Pro Gly Ile Cys Phe Glu Arg Val Asn Ser Ala Gly Asp Pro 545 550 555	1980
tat ggc aac tgt ggc aaa gtc tcg aag agt tcc ttt gcc aaa tgc gag Tyr Gly Asn Cys Gly Lys Val Ser Lys Ser Ser Phe Ala Lys Cys Glu 560 565 570	2028
atg aga gat gct aaa tgt gga aaa atc cag tgt caa gga ggt gcc agc Met Arg Asp Ala Lys Cys Gly Lys Ile Gln Cys Gln Gly Gly Ala Ser 575 580 585 590	2076
cgg cca gtc att ggt acc aat gcc gtt tcc ata gaa aca aac atc ccc Arg Pro Val Ile Gly Thr Asn Ala Val Ser Ile Glu Thr Asn Ile Pro 595 600 605	2124
ctg cag caa gga ggc cgg att ctg tgc cgg ggg acc cac gtg tac ttg Leu Gln Gln Gly Gly Arg Ile Leu Cys Arg Gly Thr His Val Tyr Leu 610 615 620	2172
ggc gat gac atg ccg gac cca ggg ctt gtg ctt gca ggc aca aag tgt Gly Asp Asp Met Pro Asp Pro Gly Leu Val Leu Ala Gly Thr Lys Cys 625 630 635	2220
gca gat gga aaa atc tgc ctg aat cgt caa tgt caa aat att agt gtc Ala Asp Gly Lys Ile Cys Leu Asn Arg Gln Cys Gln Asn Ile Ser Val 640 645 650	2268
ttt ggg gtt cac gag tgt gca atg cag tgc cac ggc aga ggg gtg tgc Phe Gly Val His Glu Cys Ala Met Gln Cys His Gly Arg Gly Val Cys 655 660 665 670	2316
aac aac agg aag aac tgc cac tgc gag gcc cac tgg gca cct ccc ttc Asn Asn Arg Lys Asn Cys His Cys Glu Ala His Trp Ala Pro Pro Phe 675 680 685	2364
tgt gac aag ttt ggc ttt gga gga agc aca gac agc ggc ccc atc cgg Cys Asp Lys Phe Gly Phe Gly Gly Ser Thr Asp Ser Gly Pro Ile Arg 690 695 700	2412
caa gca gaa gca agg cag gaa gct gca gag tcc aac agg gag cgc ggc Gln Ala Glu Ala Arg Gln Glu Ala Ala Glu Ser Asn Arg Glu Arg Gly 705 710 715	2460
cag ggc cag gag ccc gtg gga tcg cag gag cat gcg tct act gcc tca Gln Gly Gln Glu Pro Val Gly Ser Gln Glu His Ala Ser Thr Ala Ser 720 725 730	2508
ctg aca ctc atc tga gccctcccat gacatggaga ccgtgaccag tgctgctgca Leu Thr Leu Ile 735	2563
gaggagggtca cgcgtcccca aggcctcctg tgactggcag cattgactct gtggctttgc	2623
catcgtttcc atgacaacag acacaacaca gttctcgggg ctcaggaggg gaagtccagc	2683
ctaccaggca ggtctgcaga aacagtgcaa ggaagggcag cgacttcctg gttgagcttc	2743
tgctaaaaca tggacatgct tcagtgtctgc tcctgagaga gtagcagggtt accactctgg	2803
caggccccag ccctgcagca aggaggaaga ggactcaaaa gtctggcctt tcaactgagcc	2863
tccacagcag tggggggagaa gcaagggttg ggcccagtggt cccctttccc cagtgcaccc	2923
tccgccttgg cagccctgat gactgggtctc tggctgcaac ttaatgctct gatattggctt	2983
ttagcattta ttatatgaaa atagcagggt tttagttttt aatttatcag agaccctgcc	3043
accattcca tctccatcca agcaactga atggcattga aacaaactgg agaagaaggt	3103

aggagaaagg gcggtgaact ctggctcttt gctgtggaca tgcgtgacca gcagtactca 3163
 ggtttgaggg ttgcagaaa gccaggaac ccacagagtc accaaccctt catttaacaa 3223
 gtaagaatgt taaaaagtga aaacaatgta agagcctaac tccatcccc gtggccatta 3283
 ctgcataaaa atagagtgc ttttgaaata aaaaaaaaaa aaaaa 3328

<210> 4

<211> 738

<212> PRT

<213> Homo sapiens

<400> 4

Met Ala Ala Arg Pro Leu Pro Val Ser Pro Ala Arg Ala Leu Leu Leu
 1 5 10 15
 Ala Leu Ala Gly Ala Leu Leu Ala Pro Cys Glu Ala Arg Gly Val Ser
 20 25 30
 Leu Trp Asn Glu Gly Arg Ala Asp Glu Val Val Ser Ala Ser Val Arg
 35 40 45
 Ser Gly Asp Leu Trp Ile Pro Val Lys Ser Phe Asp Ser Lys Asn His
 50 55 60
 Pro Glu Val Leu Asn Ile Arg Leu Gln Arg Glu Ser Lys Glu Leu Ile
 65 70 75 80
 Ile Asn Leu Glu Arg Asn Glu Gly Leu Ile Ala Ser Ser Phe Thr Glu
 85 90 95
 Thr His Tyr Leu Gln Asp Gly Thr Asp Val Ser Leu Ala Arg Asn Tyr
 100 105 110
 Thr Val Ile Leu Gly His Cys Tyr Tyr His Gly His Val Arg Gly Tyr
 115 120 125
 Ser Asp Ser Ala Val Ser Leu Ser Thr Cys Ser Gly Leu Arg Gly Leu
 130 135 140
 Ile Val Phe Glu Asn Glu Ser Tyr Val Leu Glu Pro Met Lys Ser Ala
 145 150 155 160
 Thr Asn Arg Tyr Lys Leu Phe Pro Ala Lys Lys Leu Lys Ser Val Arg
 165 170 175
 Gly Ser Cys Gly Ser His His Asn Thr Pro Asn Leu Ala Ala Lys Asn
 180 185 190

Val Phe Pro Pro Pro Ser Gln Thr Trp Ala Arg Arg His Lys Arg Glu
195 200 205
Thr Leu Lys Ala Thr Lys Tyr Val Glu Leu Val Ile Val Ala Asp Asn
210 215 220
Arg Glu Phe Gln Arg Gln Gly Lys Asp Leu Glu Lys Val Lys Gln Arg
225 230 235 240
Leu Ile Glu Ile Ala Asn His Val Asp Lys Phe Tyr Arg Pro Leu Asn
245 250 255
Ile Arg Ile Val Leu Val Gly Val Glu Val Trp Asn Asp Met Asp Lys
260 265 270
Cys Ser Val Ser Gln Asp Pro Phe Thr Ser Leu His Glu Phe Leu Asp
275 280 285
Trp Arg Lys Met Lys Leu Leu Pro Arg Lys Ser His Asp Asn Ala Gln
290 295 300
Leu Val Ser Gly Val Tyr Phe Gln Gly Thr Thr Ile Gly Met Ala Pro
305 310 315 320
Ile Met Ser Met Cys Thr Ala Asp Gln Ser Gly Gly Ile Val Met Asp
325 330 335
His Ser Asp Asn Pro Leu Gly Ala Ala Val Thr Leu Ala His Glu Leu
340 345 350
Gly His Asn Phe Gly Met Asn His Asp Thr Leu Asp Arg Gly Cys Ser
355 360 365
Cys Gln Met Ala Val Glu Lys Gly Gly Cys Ile Met Asn Ala Ser Thr
370 375 380
Gly Tyr Pro Phe Pro Met Val Phe Ser Ser Cys Ser Arg Lys Asp Leu
385 390 395 400
Glu Thr Ser Leu Glu Lys Gly Met Gly Val Cys Leu Phe Asn Leu Pro
405 410 415
Glu Val Arg Glu Ser Phe Gly Gly Gln Lys Cys Gly Asn Arg Phe Val
420 425 430
Glu Glu Gly Glu Glu Cys Asp Cys Gly Glu Pro Glu Glu Cys Met Asn
435 440 445
Arg Cys Cys Asn Ala Thr Thr Cys Thr Leu Lys Pro Asp Ala Val Cys
450 455 460

Ala His Gly Leu Cys Cys Glu Asp Cys Gln Leu Lys Pro Ala Gly Thr
465 470 475 480

Ala Cys Arg Asp Ser Ser Asn Ser Cys Asp Leu Pro Glu Phe Cys Thr
485 490 495

Gly Ala Ser Pro His Cys Pro Ala Asn Val Tyr Leu His Asp Gly His
500 505 510

Ser Cys Gln Asp Val Asp Gly Tyr Cys Tyr Asn Gly Ile Cys Gln Thr
515 520 525

His Glu Gln Gln Cys Val Thr Leu Trp Gly Pro Gly Ala Lys Pro Ala
530 535 540

Pro Gly Ile Cys Phe Glu Arg Val Asn Ser Ala Gly Asp Pro Tyr Gly
545 550 555 560

Asn Cys Gly Lys Val Ser Lys Ser Ser Phe Ala Lys Cys Glu Met Arg
565 570 575

Asp Ala Lys Cys Gly Lys Ile Gln Cys Gln Gly Gly Ala Ser Arg Pro
580 585 590

Val Ile Gly Thr Asn Ala Val Ser Ile Glu Thr Asn Ile Pro Leu Gln
595 600 605

Gln Gly Gly Arg Ile Leu Cys Arg Gly Thr His Val Tyr Leu Gly Asp
610 615 620

Asp Met Pro Asp Pro Gly Leu Val Leu Ala Gly Thr Lys Cys Ala Asp
625 630 635 640

Gly Lys Ile Cys Leu Asn Arg Gln Cys Gln Asn Ile Ser Val Phe Gly
645 650 655

Val His Glu Cys Ala Met Gln Cys His Gly Arg Gly Val Cys Asn Asn
660 665 670

Arg Lys Asn Cys His Cys Glu Ala His Trp Ala Pro Pro Phe Cys Asp
675 680 685

Lys Phe Gly Phe Gly Gly Ser Thr Asp Ser Gly Pro Ile Arg Gln Ala
690 695 700

Glu Ala Arg Gln Glu Ala Ala Glu Ser Asn Arg Glu Arg Gly Gln Gly
705 710 715 720

Gln Glu Pro Val Gly Ser Gln Glu His Ala Ser Thr Ala Ser Leu Thr
725 730 735

Leu Ile

<210> 5
 <211> 2642
 <212> DNA
 <213> Homo sapiens

<400> 5
 taaccaaggt ttaaccatag gaattctggt gaccatcctg tgtcttcttg ctgccggatt 60
 tgtggttttat ctcaaaagga agaccttgat acgactgctg ttacaaaata agaagaccac 120
 cattgaaaaa ctaagggtgtg tgcgcccttc ccggccaccc cgtggcttcc aaccctgtca 180
 ggctcacctc ggccaccttg gaaaaggcct gatgaggaag ccgccagatt cctaccacc 240
 gaaggacaat cccaggagat tgctgcagtg tcagaatggt gacatcagca gaccctcaa 300
 cggcctgaat gtccctcagc cccagtcaac tcagcgagtg ctctctccc tccaccgggc 360
 cccacgtgca cctagcgtcc ctgccagacc cctgccagcc aagcctgcac ttaggcaggc 420
 ccaggggacc tgtaagccaa acccccctca gaagcctctg cctgcagatc ctctggccag 480
 aacaactcgg ctactcatg ccttggccag gaccccagga caatgggaga ctgggctccg 540
 cctggcacc ctcagacctg ctccacaata tccacaccaa gtgccagat ccaccacac 600
 cgcctatatt aagtgagaag ccgacacctt ttttcaacag tgaagacaga agtttgcact 660
 atctttcagc tccagttgga gttttttgta ccaactttta ggattttttt taatgtttaa 720
 aacatcatta ctataagaac tttgagctac tgccgtcagt gctgtgctgt gctatggtgc 780
 tctgtctact tgcacaggta cttgtaaatt attaatattat gcagaatggt gattacagtg 840
 cagtgcgctg tagtaggcat ttttaccatc actgagtttt ccatggcagg aaggcttggt 900
 gtgcttttag tatttttagtg aacttgaaat atcctgcttg atgggattct ggacaggatg 960
 tgtttgcttt ctgatcaagg ccttattgga aagcagtccc ccaactaccc ccagctgtgc 1020
 ttatggtacc agatgcagct caagagatcc caagtagaat ctcagttgat tttctggatt 1080
 cccatctca ggccagagcc aaggggcttc aggtccaggc tgtgtttggc tttcaggagg 1140
 gccctgtgcc ccttgacaac tggcaggcag gctcccaggg acacctggga gaaatctggc 1200
 ttctggccag gaagcttttg tgagaacctg ggttgccagac aggaatctta aggtgtagcc 1260
 acaccaggat agagactgga aactagaca agccagaact tgaccctgag ctgaccagcc 1320
 gtgagcatgt ttggaagggg tctgtagtgt cactcaaggc ggtgcttgat agaaatgcc 1380
 agcacttctt tttctcgctg tcctttctag agcactgcca ccagtagggt atttagcttg 1440
 ggaaagggtg tgtttctgta agaaacctac tgcccaggca ctgcaaaccg ccacctccct 1500
 atactgcttg gagctgagca aatcaccaca aactgtaata caatgatcct gtattcagac 1560
 agatgaggac tttcatggg accacaacta ttttcagatg tgaaccatta accagatcta 1620
 gtcaatcaag tctgtttact gcaaggttca acttattaac aattaggcag actctttatg 1680

cttgcaaaaa ctacaaccaa tggaatgtga tgttcatggg tatagtccat gtctgctatc 1740
 attattcgtg gatattggac aaagaacctt ctctatgggg catcctcttt ttccaacttg 1800
 gctgcaggaa tctttaaaag atgcttttaa cagagtctga acctatttct taaacacttg 1860
 caacctacct gttgagcatc acagaatgtg ataaggaaat caacttgctt atcaacttcc 1920
 taaatattat gagatgtggc ttgggcagca tccccttgaa ctcttcactc ttcaaatgcc 1980
 tgactaggga gccatgtttc acaaggctct taaagtgact aatggcatga gaaatacaaa 2040
 aatactcaga taaggtaaaa tgccatgatg cctctgtctt ctggactggg ttccacatta 2100
 gaagacaatt gacaacagtt acataattca ctctgagtgt tttatgagaa agccttcttt 2160
 tgggggtcaac agttttccta tgctttgaaa cagaaaaata tgtaccaaga atcttggttt 2220
 gccttccaga aaacaaaact gcatttcact ttcccgggtg tcccactgt atctaggcaa 2280
 catagtattc atgactatgg ataaactaaa cacgtgacac aaacacacac aaaagggAAC 2340
 ccagctctaa tacattccaa ctctgtatagc atgcatctgt ttattctata gttattaagt 2400
 tctttaaaat gtaaagccat gctggaaaat aatactgctg agatacatac agaattactg 2460
 taactgatta cacttggtaa ttgtactaaa gccaaacata tatatactat taaaaaggtt 2520
 tacagaattt tatgggtgcat tacgtgggca ttgtcttttt agatgcccaa atccttagat 2580
 ctggcatgtt agcccttcct ccaattataa gaggatatga accaaaaaaa aaaaaaaaaa 2640
 aa 2642

<210> 6

<211> 205

<212> PRT

<213> Homo sapiens

<400> 6

Asp Asn Gln Gly Leu Thr Ile Gly Ile Leu Val Thr Ile Leu Cys Leu
 1 5 10 15

Leu Ala Ala Gly Phe Val Val Tyr Leu Lys Arg Lys Thr Leu Ile Arg
 20 25 30

Leu Leu Phe Thr Asn Lys Lys Thr Thr Ile Glu Lys Leu Arg Cys Val
 35 40 45

Arg Pro Ser Arg Pro Pro Arg Gly Phe Gln Pro Cys Gln Ala His Leu
 50 55 60

Gly His Leu Gly Lys Gly Leu Met Arg Lys Pro Pro Asp Ser Tyr Pro
 65 70 75 80

Pro Lys Asp Asn Pro Arg Arg Leu Leu Gln Cys Gln Asn Val Asp Ile
 page 17

85

90

95

Ser Arg Pro Leu Asn Gly Leu Asn Val Pro Gln Pro Gln Ser Thr Gln
 100 105 110

Arg Val Leu Pro Pro Leu His Arg Ala Pro Arg Ala Pro Ser Val Pro
 115 120 125

Ala Arg Pro Leu Pro Ala Lys Pro Ala Leu Arg Gln Ala Gln Gly Thr
 130 135 140

Cys Lys Pro Asn Pro Pro Gln Lys Pro Leu Pro Ala Asp Pro Leu Ala
 145 150 155 160

Arg Thr Thr Arg Leu Thr His Ala Leu Ala Arg Thr Pro Gly Gln Trp
 165 170 175

Glu Thr Gly Leu Arg Leu Ala Pro Leu Arg Pro Ala Pro Gln Tyr Pro
 180 185 190

His Gln Val Pro Arg Ser Thr His Thr Ala Tyr Ile Lys
 195 200 205

<210> 7

<211> 908

<212> DNA

<213> Homo sapiens

<400> 7

agcaaggcag gaagctgcag agtccaacag ggagcgcggc cagggccagg agcccgtggg 60
 atcgcaggag catgctgcta ctgcctcact gacactcatc tgagccctcc catgacatgg 120
 agaccgtgac cagtgtgtgt gcagaggagg tcacgcgtcc ccaaggcctc ctgtgactgg 180
 cagcattgac tctgtggctt tgccatcgtt tccatgacaa cagacacaac acagttctcg 240
 gggctcagga ggggaagtcc agcctaccag gcaggtctgc agaaacagtg caaggaaggg 300
 cagcgacttc ctggttgagc ttctgctaaa acatggacat gcttcagtgc tgctcctgag 360
 agagtagcag gttaccactc tggcaggccc cagccctgca gcaaggagga agaggactca 420
 aaagtctggc ctttactga gcctccacag cagtggggga gaagcaaggg ttgggcccag 480
 tgtccccttt cccagtgac acctccgcct tggcagccct gatgactggt ctctggctgc 540
 aacttaatgc tctgatatgg ctttttagcat ttattatatg aaaatagcag ggtttttagtt 600
 ttttaatttat cagagaccct gccacccatt ccattctccat ccaagcaaac tgaatggcat 660
 tgaaacaaac tggagaagaa ggtaggagaa agggcgggtga actctggctc tttgctgtgg 720
 acatgcgtga ccagcagtac tcaggtttga gggtttgag aaagccaggg aaccacaga 780
 gtcaccaacc cttcatttaa caagtaagaa tggttaaaaag tgaaaacaat gtaagagcct 840

▲ aactccatcc cccgtggcca ttactgcata aaaatagagt gcattttgaa ataaaaaaaa 900
 aaaaaaaaa 908

<210> 8
 <211> 34
 <212> PRT
 <213> Homo sapiens

<400> 8
 Glu Ala Arg Gln Glu Ala Ala Glu Ser Asn Arg Glu Arg Gly Gln Gly
 1 5 10 15
 Gln Glu Pro Val Gly Ser Gln Glu His Ala Ser Thr Ala Ser Leu Thr
 20 25 30

Leu Ile

<210> 9
 <211> 23
 <212> DNA
 <213> Homo sapiens

<400> 9
 cctgtgacct cccagagttc tgc 23

<210> 10
 <211> 23
 <212> DNA
 <213> Homo sapiens

<400> 10
 gaattcctat ggttaaacct tgg 23

<210> 11
 <211> 24
 <212> DNA
 <213> Homo sapiens

<400> 11

cttgctgccg gatttgtggt ttat 24

<210> 12

<211> 24

<212> DNA

<213> Homo sapiens

<400> 12
agttgactgg ggctgagggga catt 24

<210> 13

<211> 23

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (15)..(15)

<223> Mismatch (g instead of c)

<400> 13
ggactctgca gcttgctgcc ttg 23

<210> 14

<211> 23

<212> DNA

<213> Homo sapiens

<400> 14
ggactctgca gcttcctgcc ttg 23

<210> 15

<211> 21

<212> DNA

<213> Homo sapiens

<400> 15
tcggggctca ggaggggaag t 21

<210> 16
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <400> 16
 ggctgccaa ggcggaggtgt c 21

 <210> 17
 <211> 24
 <212> DNA
 <213> Homo sapiens

 <400> 17
 ggccgggtga tctttggtct ctct 24

 <210> 18
 <211> 24
 <212> DNA
 <213> Homo sapiens

 <400> 18
 cccggctgtc tgtcttggtg ctct 24

 <210> 19
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <400> 19
 ctgctgagat gggaaagggc t 21

 <210> 20
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <400> 20
 ttcaggataa tcacctgagc a 21

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.